



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ



I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

I also certify that the attached copy of the request for grant of a Patent (Form 1/77) bears an amendment, effected by this office, following a request by the applicant and agreed to by the Comptroller-General.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.


Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed

Dated 15 December 2000





The Patent Office  
14 APR 1998

1/77  
15APR98 E353090-3 D00019  
P01/7700 25.00 - 9807922.1

**Request for grant of a patent**

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

**9807922.1**

The Patent Office

Cardiff Road  
Newport  
Gwent NP9 1RH

1. Your reference **P19015GB** **G19859P/HGH**

2. Patent application number  
(The Patent Office will fill in this part)

3. Full name, address and postcode of the or of each applicant (underline all surnames)

**COBRA THERAPEUTICS LIMITED**  
**The Science Park**  
**University of Keele**  
**Keele**  
**Staffordshire**  
**ST5 5SP**

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

**United Kingdom**

**0747108009**

4. Title of the invention

**Purification of cellular components that are substantially RNase free**

5. Name of your agent (if you have one)

**Carpmaels & Ransford**

**6/12/00 Form S1/77**

**Harrison Gaddard**  
**Forster**

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

**43 Bloomsbury Square**  
**London**  
**WC1A 2RA**

**Tower House**

**Merton Way**

**LEEDS**

Patents ADP number (if you know it)

**83001**

**LS2 8PA**

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number  
(if you know it)

Date of filing  
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

- a) any applicant named in part 3 is not an inventor, or
- b) there is an inventor who is not named as an applicant, or
- c) any named applicant is a corporate body

**Yes**

See note (d))

# Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description	57
Claim(s)	3
Abstract	1
Drawing(s)	4 + 4



10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Yes /

Request for substantive examination (*Patents Form 10/77*)

Any other documents  
(please specify)

11. I/We request the grant of a patent on the basis of this application.

Signature

Date



14th April 1998

12. Name and daytime telephone number of person to contact in the United Kingdom

Huw G. Hallybone  
Carpmaels & Ransford

0171 242 8692

## Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

## Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

**PURIFICATION OF CELLULAR COMPONENTS  
THAT ARE SUBSTANTIALLY RNase FREE**

Field of the Invention

The invention relates to the production of RNA free cellular components, and a method  
5 of removing RNA from preparations of cellular components.

Background of the Invention

The ribonuclease enzyme family (hereafter referred to as RNases) has been extensively studied. Numerous RNases have been characterized and the genes encoding some of these proteins have been cloned.

10 RNases hydrolyze one or more of the phosphodiester bonds in single stranded and double stranded RNA as well as RNA in RNA:DNA hybrids. RNases differ in their specificity for either a particular form of the RNA substrate (for example single stranded, double stranded or in a DNA:RNA hybrid) or their specific point of RNA cleavage.

15 One biological activity of RNase enzymes is the processing of mature molecules of RNA from precursor forms (Genes, Benjamin Lewin ed., John Wiley & Sons, 2nd edition, p.395, 1985). Certain RNases can affect the growth and differentiation of mammalian cells by virtue of an intrinsic anti-tumor activity (references within Ribo *et al.* Prot. Express. And Purif. 7: 253-261, 1996).

20 RNA is a major contaminant of preparations from cell lysates. For example, plasmid DNA preparations contain RNA that is difficult to remove by anion exchange or size exclusion chromatography because it is similar in size and charge to DNA. RNase A hydrolyzes RNA after C and U residues by cleaving between the 3'-phosphate group of a pyrimidine ribonucleotide and the 5'-hydroxyl of the adjacent nucleotide. This  
25 enzyme is used to degrade RNA to low molecular weight species that no longer copurify with plasmid or genomic DNA. RNase I can also be used to remove RNA from preparations of plasmid or genomic DNA. RNase A is also commonly used for the enzymatic digestion of host derived RNA during the production of recombinant protein from *E. coli*.

A prior art method of removing RNA from a sample is to add a large quantity of an exogenously produced RNase. For example, to remove RNA from plasmid DNA bovine RNase A is added to a final concentration of 100µg/ml (Qiagen Plasmid Handbook February 1995, Qiagen Ltd. Unit 1 Tillingbourne Court, Dorking Business  
5 Park, Dorking, Surrey RH4 1HJ, UK). Bovine RNase A is commonly used at a final concentration of approximately 10-100µg/ml for the enzymatic digestion of host derived RNA during the production of recombinant protein from *E. coli*. A chief disadvantage of prior art methods is that the cellular component that is treated according to these methods often contains residual RNase.

- 10 There are limitations to using exogenously produced RNases. First, it is difficult to purify large amounts of RNase from the tissue of origin. This is presumably because high concentrations of active RNase will degrade host cell RNA and impair normal cell functions to a level that can be toxic to a cell. Therefore, it is difficult to produce large quantities of active RNase by expression in cells because high concentrations of active  
15 RNase will be toxic to a cell. It is also difficult to produce large quantities of active RNase by expression in cells because RNase can be sensitive to proteases of the host in which it is being synthesized, and because RNases that are over-produced and accumulate as in e.g. *E. coli* inclusion bodies cannot always be correctly refolded. The most significant limitation to using exogenously produced RNase is that, if the  
20 exogenously added RNase is of animal origin, following RNase treatment the presence of residual enzyme can contaminate a DNA preparation, thereby rendering it unacceptable for certain applications, including gene therapy. Third, it is expensive to produce RNase in large quantities.

There is a need in the art for plasmid DNA and protein preparations that are  
25 substantially free from contaminating RNA.

There is a need in the art for methods of producing RNA-free cellular components that are suitable for administration to human subjects.

There is a need in the art for a method of removing RNA from cellular components that does not rely on incubating a cellular lysate or purified component with added RNase.

30 Summary of the Invention

The invention features a method of preparing a substantially RNA-free cellular component comprising culturing cells producing the cellular component in a medium and lysing said cells to produce a cell lysate, wherein said cell lysate contains said cellular component and sufficient RNase activity to degrade substantially all of the RNA molecules present in said cell lysate.

In a preferred embodiment, the RNase is produced by cells producing the cellular component.

Alternatively, the RNase is produced by cells in the medium other than those cells producing the cellular component.

The invention also features a method of preparing a substantially RNA-free cellular component, comprising culturing and lysing cells producing the cellular component and cells producing an RNase in an amount sufficient to degrade substantially all of the RNA present in the preparation.

Preferably, the cells producing the cellular component also produce the RNase and the culture and lysate contain cells producing the cellular component and an RNase in an amount sufficient to degrade substantially all of the RNA present in the preparation.

Preferably, the cellular component and the RNase are not produced by the same cell.

In preferred embodiments of both of the above-described inventive methods, the cellular component is one of a DNA, a protein, and a carbohydrate. Preferably, the cellular component is one of a recombinant DNA, a recombinant protein and a recombinant carbohydrate.

In preferred embodiments of both of the above-described inventive methods the gene encoding said RNase is integrated into the genome of the cell producing the RNase.

In some methods of the invention, it is preferred that the RNase is non-specific. Such a non-specific RNase may be RNase A, RNase M or RNase I.

In preferred embodiments of both of the above-described inventive methods, a cell producing an RNase produces the RNase in a regulated manner.

Preferably, the RNase produced by the host cell is overproduced, either by an inducible production, or by constitutively production. The RNase overproduced by the host cell

also may be secreted out of the host cell cytoplasm, for example, secreted into the host cell periplasm or secreted out of the host cell into the medium.

In some methods, it is preferred that the RNase is a non-specific RNase.

5 The invention also features a composition comprising a host cell that produces a recombinant DNA, a recombinant protein, or a recombinant carbohydrate and also produces an RNase in a regulated manner.

RNase produced in said regulated manner is overproduced, or inducibly overproduced, or constitutively overproduced. The RNase produced by the host cell also may be secreted out of the host cell cytoplasm, for example, secreted into the host cell  
10 periplasm or secreted out of the host cell into the medium.

In some methods, it is preferred that the RNase is a non-specific RNase.

The invention also features a composition comprising a host cell that produces a recombinant DNA, a recombinant protein, or a recombinant carbohydrate and a host cell that produces an RNase in a regulated manner.

15 RNase produced in said regulated manner is overproduced, or inducibly overproduced, or constitutively overproduced. The RNase produced by the host cell also may be secreted out of the host cell cytoplasm, for example, secreted into the host cell periplasm or secreted out of the host cell into the medium.

The invention also features a pharmaceutical composition comprising a cellular  
20 component that is substantially RNA-free, in a pharmaceutically acceptable carrier. The present invention also provides a pharmaceutical composition comprising a cellular component that is substantially RNA-free obtainable by the methods of the present invention, in a pharmaceutically acceptable carrier.

25 Preferably, the cellular component that is substantially RNA-free is made according the methods described herein.

As used herein, "substantially RNA-free" and "substantially all of the RNA molecules" refers to the exceedingly small amount of RNA that is permitted in a preparation of a cellular component that is useful for administration to humans. Acceptable exceedingly low levels of RNA are as follows.



"Substantially RNA-free DNA" refers to the presence in a sample containing the cellular component of less than 1%, preferably less than 0.2% and most preferably less than 0.1%-0.01% (w/w of RNA/DNA in the sample).

"Substantially RNA-free protein" refers to a protein preparation that contains less than 1% and preferably less than 0.2% and most preferably less than 0.1%-0.01% (w/w) of RNA/protein. When the protein is a therapeutic protein "substantially RNA-free protein" refers to a therapeutic protein preparation that contains less than 500 pg of RNA/dose, preferably less than 100 pg of RNA/dose, most preferably less than 5-10 pg of RNA/dose.

10 "Substantially RNA-free carbohydrate" refers to a carbohydrate preparation that contains less than 1% and preferably less than 0.2% and most preferably less than 0.1%-0.01% (w/w) of RNA/carbohydrate.

Preferably, the cell lysate will contain significantly less RNase protein than prior art methods where exogenous RNase is added to cell lysate. "Significantly less" in this context refers to less than 10µg - 50 µg RNA/ml lysate, preferably less than 10µg RNA/ml of lysate and more preferably less than 1µg RNA/ml.

As used herein, "cellular component" refers to any one of DNA, including but not limited to plasmid DNA, cosmid DNA, yac DNA, episomal DNA, or genomic DNA; recombinant protein, and recombinant carbohydrate.

20 As used herein, "recombinant" refers to DNA, protein or carbohydrate that is not naturally occurring in a cell or produced in an amount that is not normally produced by the cell.

As used herein, "cell" refers to a gram negative or gram positive bacteria, including but not limited to *E. coli*, *Salmonella typhimurium*, *Bacillus spp.*, *Streptomyces spp.* and *Pseudomonas aeruginosa*.

"Regulated manner" refers to gene expression, for example, of an RNase gene in a host cell that is transcriptionally regulated, for example, constitutively or inducibly. It also refers to regulation at the level of protein production, for example, the use of a signal sequence or a fusion protein to direct a protein to the host cell periplasm or out of the host cell and into the medium surrounding the host cell.

“Inducible” or “induction” refers to transcriptional activation or derepression; “constitutive” refers to continuous transcription often at a constant level; “signal sequence” refers to an amino acid sequence that directs secretion of the protein into the periplasm or out of the cell; “secretion” refers to movement of a secreted protein out of the host cell cytoplasm and into the periplasm or host cell culture medium; “periplasm” refers to the host cell compartment between inner and outer cell membranes of, for example, *E. coli*.

“Overexpresses” or “overproduces” refers to gene expression at a level higher than that normally expressed in the host cell; “overproduced” refers to production in an amount higher than that normally produced by the cell. Thus, over-expression of a given gene or overproduction of a gene product or a recombinant DNA encompasses production of 10% or more, 50% or more, 100% or more, or even up to over 200-500% more of the gene product than the cell normally produces in the absence of the over-expressed gene.

The invention provides DNA, protein or carbohydrate preparations that are substantially free of contaminating RNA. The invention is particularly useful in providing preparations that involve administration to humans, but also for other uses, including for accurate determination of DNA concentrations by absorbance at 260nm (where RNA also absorbs), and for investigating both DNA/RNA and DNA/protein interactions. Substantially RNA-free protein is required for therapeutic applications, for estimating protein concentrations by absorbance at 280nm, for analysis of protein crystal structures, for studying protein/DNA and protein/RNA interactions, and for certain protein purification protocols where nucleic acid binding reduces the affinity of protein for ion-exchangers or reduces the capacity of ion-exchangers for protein.

The invention is applicable to any use requiring a preparation of a cellular component, such as DNA, protein, carbohydrate, that is substantially free of contaminating RNA, for example, for preparation of a cellular component such as DNA or a protein or a carbohydrate that is suitable for therapeutic use in humans.

Further features and advantages of the invention will become more fully apparent in the following description of the embodiments and drawings thereof, and from the claims.

#### BRIEF DESCRIPTION OF DRAWINGS

The drawings are briefly described.

Figure 1 shows plasmid pN1.

Figure 2 is a diagrammatic representation of the strategy for introducing and deleting plasmid DNA into the *E. coli* chromosome and for deleting chromosomal DNA from the *E. coli* chromosome by insertional mutagenesis and homologous recombination.

Figure 3 is an LB agar plate assay demonstrating expression of RNase I from TOP10F cells transformed with pCR3.1RNase I.

Figure 4 is an LB agar plate assay demonstrating IPTG inducible expression from DH5 $\alpha$  cells transformed with pUC18RNase I.

Figure 5 is an LB agar plate assay demonstrating the sensitivity of purified pUC18 RNase I to alkaline lysis.

Figure 6 is an ethidium bromide stained agarose gel demonstrating the affects of lysis conditions on the activity of pUC18 RNase I.

#### DESCRIPTION

The invention provides for a method of producing substantially RNA-free cellular components that includes producing a selected cellular component and an RNase in the same host cell or different host cells that are co-cultured such that, upon lysis of the cell(s), sufficient RNase activity is present so as to degrade substantially all of the RNA molecules in the lysate.

The cellular component may be a nucleic acid, protein or carbohydrate that is produced according to the invention substantially RNA-free. This may be achieved, for example, using a gene encoding a cellular component that is introduced into a host cell that has been engineered to express an RNase protein, or by culturing and lysing two different host cells that individually produce the cellular component and the RNase.

#### 25 RNase Genes and RNases Useful According to the Invention

Genes encoding a variety of RNase proteins have been cloned. Ideally, the RNase of the invention is non-specific (for example, bovine RNase A and *E. coli* RNase M used at high concentrations) and therefore will degrade all types of RNA molecules. Non-specific RNases hydrolyze phosphodiester bonds in single stranded or double stranded RNA or RNA/DNA molecules, anywhere along the length of the RNA molecule.

The protein products and the genes for bovine RNase A (Tarragona-Fiol *et al.*, Gene, 118: 239-245, 1992, Schein *et al.*, Biochem. J., 283: 137-144, 1992, Laity *et al.*, PNAS 90: 615-619, 1993, delCardayre, Prot. Engineer. 8: 261-273, 1995, Okorokov *et al.*, Prot. Express. Purif. 6: 472-480), and murine and rat RNase A homologs (Schein *et al.*,  
 5 supra) have been cloned and characterized.

RNase H hydrolyzes RNA molecules that are hybridized to a complementary DNA strand and is used in molecular biology to degrade the RNA strand after first-strand synthesis in the production of double-stranded cDNA. RNase H is also used to remove polyA tails from mRNA that has hybridized to oligo-dT<sub>12-18</sub> (described in U.S. Patent  
 10 No.: 5,459,055).

RNase I cleaves the phosphodiester bond of single stranded RNA 3' of any ribonucleoside and is used for RNase protection assays, mapping and quantitation of RNA and mismatch detection. *E. coli* RNase I and RNase M (Meador and Kennell, 1990, Ribo *et al.*, Prot. Express. Purif. 7: 253-261, 1996) have been cloned, isolated and  
 15 characterized.

RNase T1 preferentially cleaves 3' to G residues and is used for RNA sequence analysis and RNA fingerprinting. *Aspergillus oryzae* RNase T1 and T2 (Fujimura *et al.*, FEBS, 265:71-74, 1990, Quass *et al.*, Eur. J. Biochem. 173: 617-622, 1988) have been cloned, isolated and characterized.

20 RNase4 (Seno *et al.*, Biochim. Et Biophys. Acta, 1261: 424-426-1995) has been cloned, isolated and characterized.

Human pancreatic RNase (Russo *et al.*, FEBS 333: 233-237, 1993) has been cloned, isolated and characterized.

*Bacillus amyloliquefaciens* BaRNase (Hartley, J. Mol. Biol. 202: 913-915, 1988) has  
 25 been characterized with respect to synthesis and processing.

#### Expression of a Selected RNase gene in a Host Cell

According to the invention, an RNase gene can be modified so that the RNase protein is expressed in either the periplasmic space or the cellular supernatant, for example, by directing the location of the RNase using a signal sequence or by creating a fusion  
 30 protein that is secreted.

Alternatively, expression of the RNase may be selectively controlled by placing the RNase gene under the control of an inducible promoter.

Alternatively, the RNase gene may be introduced into a host cell and integrated into the chromosome of the host cell.

- 5 Vectors used to overproduce RNase have been modified to minimize problems due to toxic effects of improperly folded RNase on the host cell in which the RNase is being produced, sensitivity of unfolded RNase to proteases present in the host cell and the improper refolding of some RNases that have been produced in inclusion bodies.

#### Integration Of An RNase Gene Into The Chromosome Of The *E. coli* Host Strain

- 10 The following methods may be used to provide a host cell containing an RNase gene integrated into the host cell chromosome.

##### 1. Cloning RNase A into pN1

The RNase A gene was isolated from the expression vector pQR162 (Tarragona-Fiol *et al.*, supra) by PCR (Current Protocols in Molecular Biology, Ausubel, F. M *et al.* eds.,

- 15 John Wiley & Sons Inc.) using the primers.

5'-CTCGAATTCAATGTTCTTGGAGGATGATTG-3'

5'-TACGAATTCGGCCTTAGGTAGAGACCTAC-3'.

- 20 The isolated RNase gene was inserted into the EcoRI site of pUC18 such that an inframe fusion was created between the lacZ gene and the hexapeptide located 5' to the RNase A gene. The expression of the RNase A gene and the hexapeptide was controlled by the lacZ promoter. The LacZ-RNase A construct was then excised by a Hae II digest, blunt ended, and inserted into pN1 (Figure 1) previously digested with Sty I and blunt ended.

The resulting construct is referred to as pN1RNase A.

- 25 2. Introduction of pN1RNase into *E. coli*

The resulting construct, referred to as pN1RNase A, consists of an RNase expression cassette flanked on both sides by the *E. coli* dif locus containing chromosomal DNA homology. The dif locus is a 28 base pair region that comprises the recognition site for

XerC and XerD recombinases (Hayes and Sherratt, J. Mol. Biol., 266: 525- 537, 1997, Kuempel, *et al.*, The New Biologist 3: 799-811, 1991).

The resultant pN1RNase A construct was used, to shuttle the RNase A gene cassette into the *E. coli* strains JC7623 and then DH1 by the following method.

- 5 pN1RNase A was linearized with Sal I (or another suitable restriction enzyme), or subjected to a partial digest reaction with Sal I. The resulting fragment (containing the excised but intact RNase expression cassette) was used to transform calcium-competent JC7623 cells (recB21, sbcC201, recC22, sbcB15; Horii and Clark, J. Mol. Biol. 80: 327-344, 1973, Lloyd and Buckman, J. Bacteriol. 164: 836-844, 1985) by linear  
10 transformation as described by Winans *et al.*, J. Bacteriol. 161: 1291-1221, 1985 and Jasin and Schimmel, J. Bacteriol. 159: 783-786, 1984.

- P1 phage transduction was then used to move this chromosomal construct into the dif locus of *E. coli* DH1 (F-, *supE44*, *recA1*, *endA1*, *gyrA96*, *thi*-, *hsdr17*, *relA1* as described in Hanahan, Mol. Biol. 166: 557-580, 1983, and Bachmann, in *Escherichia coli* and *Salmonella typhimurium*, Cellular and Molecular Biology, ed. Neidhardt, F. C.  
15 *Et al.*, ASM, pp. 1190-1219, 1987) or another suitable DNA vector host strain as detailed in Figure 2. To facilitate integration of the RNase gene into the host cell chromosome, the target host strain was first rendered transiently RecA+ by transformation with the unstable, *recA*- containing plasmid pPE13 (Hickson *et al.*,  
20 *Escherichia coli* Mol. Gen. Genet., 184: 68-72, 1981). Transformation of the host strain with pPE13 was performed using the calcium phosphate method of preparing competent cells, as described in Short Protocols in Molecular Biology, Ausubel *et al.*, eds, Wiley, 3rd edition, 1993).

### 3. PCR Amplification of pN1RNase I

- 25 The RNase I gene was amplified from the *E. coli* strain DH1 by PCR using the primers, 5' GGTCCTGGGGTGATTATTTACGGCTGTGGC-3', and 5'-GTTTAACTCACATGATGATACTGACTGTTG-3', and cloned into the TA cloning vector pCR3.1 (Invitrogen) to generate pCR3.1RNase I.

- The RNase I gene was then amplified from pCR3.1RNase I by PCR using the primers,  
30 5'-TCCAGAATTCCATGAAAGCATTCTGGGG-3' and 5'-GTTGAATTCACATGATGATACTGACTGTTG-3', (according to the method of

Ausubel *et al.*, Current Protocols, supra) and cloned into the *EcoRI* site of pUC18 such that the RNase I gene was fused in frame to the remaining sequences of the *lacZ* gene.

The resulting plasmid comprised an RNase I gene under the control of the *lacZ* promoter wherein the expression of the RNase I gene was increased in the presence of 0.5mM IPTG. The LacZ-RNase I construct was then excised by a Hae II digest, blunt ended, and inserted into pN1 which had been digested with Sty I and blunt ended. The resulting pN1RNaseI construct was used, as above, to shuttle the RNaseI gene cassette into the *E. coli* strains JC7623 and then DH1, as above.

#### Integration Of An RNase Gene Into The Chromosome Of Other Cell Types

##### 10 1. Production of Recombinant nucleic acids in other cell types

###### 1. *S. typhimurium*.

An RNase gene may be inserted into the chromosome of *S. typhimurium* as described in Bachmann *et al.*, 1977, in *E. coli* and *Salmonella typhimurium*, Cellular and Molecular Biology, ed. Neidhardt *et al.*, ASM, or as described in Nakayama *et al.*, 15 Bio/Technology, 6: 693-697, 1988).

##### 2. Production of Recombinant protein in other cell types

###### 1. *Bacillus spp.*

An RNase gene may be inserted into the chromosome of *Bacillus spp.* as described in Solma *et al.*, J. Bact., 173: 6889-6895.

##### 20 2. *Streptomyces spp.*

An RNase gene may be inserted into the chromosome of *Streptomyces spp.* as described in Chpt 6 in Plasmids A practical Approach (2nd edition) Edited by K.G.Hardy or as described in Motamedi *et al.*, Gene, 160:25-31, 1995.

Integrative expression vectors for heterologous expression of genes in *Streptomyces* 25 have been developed. The vectors are comprised of a strong constitutive promoter, P<sub>E</sub>, a synthetic ribosome-binding site, ATG start codon, multiple cloning site, transcription terminator and hygromycin-resistance-encoding gene (Hy<sup>R</sup>). The vectors also contain a ColE1 replicon for propagation in *Escherichia coli* and a wide-host-range *Streptomyces* integration element, the mini-circle, to direct the insertion of the vectors into the 30 *Streptomyces* genome at the mini-circle attachment site. Hy<sup>R</sup> transformants are stable in

the absence of drug selection. Conjugative derivatives can be constructed by incorporating *oriT*, the origin of transfer of the IncP plasmid RK2, into these vectors, and conjugal transfer from an appropriate *E. coli* donor to *Streptomyces lividans* (*Sl*) can be done. Derivatives of these vectors potentially useful for gene disruption, as well as complementation, have also been described. It has been demonstrated that replicative forms of the constructed mini-circle-based vectors in *Sl*, can co-exist with the integrated copy of the vector, without any apparent instability problems. The utility of these vectors has been demonstrated by expression of the gene encoding 31-O-methyltransferase, which is involved in methylation at position 31 of the immunosuppressive drug FK506. in *Sl* (Motamedi *et al.*, Supra).

### 3. *Pseudomonas* spp.

An RNase gene may be inserted into the chromosome of *Pseudomonas* spp. as described in Gene replacement in *Pseudomonas aeruginosa* in Methods in Enzymology, 235: 466-474 1994, Molecular Microbiology, 6:1195-1204, 1992 or as described in Schweizer and Hoang, Gene, 158: 15-22, 1995).

A novel pUC19-based gene replacement vector has been developed. This vector incorporates (i) the counterselectable *sacB* marker, (ii) a *lacZα* allele for blue-white screening, (iii) an *oriT* for conjugation-mediated plasmid transfer and (iv) unique cloning sites for *Sma*I and the rare-cutting meganuclease I-*Sce*I. These rare restriction sites are also present on the helper plasmid pUC19Sce. The replacement vector is engineered to contain few restriction sites to gain greater access to restriction sites within cloned DNA fragments, thus facilitating their genetic manipulation. The usefulness of the system was demonstrated by chromosomal integration of a newly constructed *xylE::Gm<sup>R</sup>* fusion cassette into the *glpD* gene of *Pseudomonas aeruginosa* (Schweizer and Hoang, supra).

### Preparation of Cells Expressing RNase and Recombinant DNA Or Protein

Cells which express an RNase gene and a DNA or protein of interest are prepared as follows.

*E. coli* host strains DH1, DH5α, DH10B, JM109, DL795, XL1-Blue and TG1 can be used for production of recombinant plasmid DNA, and *E. coli* strains BL21(DE3), JM109(DE3) and HB101 can be used for the production of recombinant protein.



### 1. Curing *E. coli* of Rec A containing plasmid

Following integration of the heterologous RNase expression cassette into the chromosome of the DNA vector host strain as described in the section entitled introduction of pN1RNase into *E. coli*, the host cells will be 'cured' of the recA-containing plasmid pPE13 by one of the following methods: serial subculture in the absence of the selectable marker drug encoded for by the plasmid (e.g. selection in the absence of ampicillin), titration with DNA intercalators such as ethidium bromide, or growth in 10% SDS (Tolmasky *et al.*, Virulence plasmids. In Plasmids: a practical approach, Ed. K.G. Hardy, IRL Press Oxford, UK, 1993). Plasmid free segregants will be selected by replica plating (plus and minus selection), and 'curing' will be confirmed by a rapid procedure for the screening of recombinant plasmid DNA (Birnboim and Doly, Nucleic Acids Res. 7: 1513-1523, 1979).

### 2. Transformation of *E. coli* with recombinant plasmid

Once cured the host cells will be made competent for DNA transformation, transformed with the recombinant plasmid (Sheen, p.1.8.5, in Ausubel, Current Protocols, supra) and selected under the appropriate conditions.

### 3. Purification of Recombinant Nucleic Acid or Protein

#### Recombinant plasmid DNA

For production of recombinant plasmid DNA, the transformed clones will be used directly to propagate plasmid DNA. Plasmid DNA is prepared according to procedures well-known in the art. See, for example, Birnboim and Doly, supra, Tolmasky *et al.*, supra, and Ausubel, Current Protocols, supra).

Plasmid DNA can be prepared as described in WO 97/29190 except that the step of adding an exogenous Rnase is omitted. Alternatively, the plasmid DNA can be prepared according to the method of Horn *et al.*, (Human Gene Therapy, 6: 565-573, 1995).

#### Production of bacterial culture

Fermentation in complete TB medium containing the appropriate concentration of antibiotic for selection (e.g. 50 µg/ml kanamycin for kan selected plasmids) will be

performed in a 10-liter Braun Biostat ED fermenter. Fermentation conditions will be maintained as follows: Temperature will be controlled at 30°C, agitation at 600 rpm, airflow at 10 liters/min, and pH at 7.0. Bacteria will be harvested in late log phase (10-11 hr post inoculation) at a final OD<sub>600</sub> of approximately 30 in a Jouan centrifuge at 3,000 x g for 30 min and stored at -20°C. A 0.75-ml aliquot will be withdrawn at harvest and processed for miniprep analyses (Promega Wizard, Madison, WI). According to this method, the average fermentation should yield 4 mg of plasmid DNA/liter (Horn *et al.*, supra).

#### Lysis and plasmid recovery

10 Fresh or frozen bacteria will be lysed using a modification of the standard alkaline method. The yield should be equivalent from fresh and frozen cell paste. Approximately 180 grams of bacteria will be resuspended in 1.4 liters of 61 mM glucose, 10mM Tris, 50 nM EDTA pH 8.0 buffer prechilled to 4-10°C. Lysis will occur after the addition of 2.8 liters of 0.2 N NaOH, 1% SDS buffer prechilled to 4-10°C  
15 followed by incubation on ice for 10 min. Cellular debris will be precipitated with the addition of 2.1 liters of 3 M potassium acetate pH 5.0 prechilled to 4-10°C, followed by incubation on ice for 10 min. Cellular debris in the lysate will be removed by filtration through Miracloth (Calbiochem, San Diego, CA). Once the cell debris has been removed, the lysate will be poured twice through a minimum of 16 layers of Miracloth  
20 each pass. Plasmid DNA will be precipitated from the cleared lysate by the addition of 0.6 volumes of -20°C 2-propanol followed by incubation for 1-2 hr at room temperature. Approximately 4.6 liters of the clear liquid layer above the enriched plasmid precipitate will be decanted. The concentrated plasmid DNA precipitate at the bottom of the vessel will be transferred to 250-ml centrifuge bottles and centrifuged at  
25 10,000 x g for 30 min in a GSA rotor and Sorvall RC5B centrifuge. Supernatants from the centrifugation will be discarded and the pellets will be drained and resuspended in approximately 200 ml of 10 mM Tris, 1 mM EDTA pH 8.0 (TE buffer). Solid ammonium acetate will be dissolved in the plasmid DNA solution with stirring to a final concentration of 2.5 M. The ammonium acetate solution containing the plasmid DNA  
30 will then be incubated on ice for 15 min. The resulting precipitate was removed by centrifuging at 10,000 x g for 20 min as described above. The supernatant will be filtered through a 0.8 µm nitrocellulose disposable membrane filter bottle unit (The

Nalge Company, Rochester, NY) and sufficient 30% PEG-8000 in 1.6 M NaCl will be added to achieve a final concentration of 10% PEG-8000 (Lis and Schleif, 1975; Lis, 1980). Plasmid DNA will be allowed to precipitate from this solution for 8-24 hr at 4°C. Plasmid DNA will be collected by centrifuging at 10,000 x g for 30 min as described above. The pellet containing the plasmid will be resuspended in 10 ml of column buffer (10 mM Tris, 1 mM EDTA, 150 mM NaCl, pH 8.0). The plasmid will then be filtered and applied directly to the column as described below.

#### Pharmacia Sephacryl S-1000 superfine gel filtration chromatography

Two Pharmacia XK26/100 columns will be pressure-packed, using a Pharmacia FPLC system according to the manufacture's instructions for this support, to final bed heights of approximately 85 cm each, resulting in a total column volume of approximately 900 ml. The running buffer will be 10 mM Tris, 1 mM EDTA, 150 mM NaCl, pH 8.0. The flow rate will be 0.75 ml/min. Partially purified plasmid DNA from the PEG-8000 precipitation described above will be passed through a 0.1- to 0.22- $\mu$ m sterile cellulose acetate Acrodisc syringe filter (Gelman, Ann Arbor, MI) and loaded onto the column. Column operation and fraction collection will be automated with a Pharmacia FPLC. Five-milliliter fractions will be collected in 15-ml disposable conical tubes. Following chromatography, the column and FPLC will be sanitized with 2 column volumes of 0.2 M NaOH. Fractions will be analyzed with 0.8% agarose gel electrophoresis, and fractions containing predominantly supercoiled plasmid DNA will be pooled in sterile disposable centrifuge bottles; the pool will be precipitated with two volumes of cold ethanol.

#### Production of sterile bulk plasmid DNA

The ethanol-precipitated column-purified plasmid DNA will be collected by centrifugation at 10,000 x g for 30 min. The plasmid pellet will be drained and air-dried under aseptic conditions in a laminar flow hood. The dried pellets will be resuspended in Lactated Ringer's (Baxter, Deerfield, IL) and diluted to an appropriate concentration for the intended dosage. The resulting solution will be subjected to a sterilizing filtration through a pyrogen-free 0.2- $\mu$ m Acrodisc filter (Gelman, Ann Arbor, MI). The sterile bulk plasmid DNA will be aliquoted and stored frozen at -70°C (Horn *et al.*, *supra*).

### Recombinant Protein

For the production of recombinant protein, the transformed plasmid will comprise an expression cassette for the production (overexpression) of a recombinant (usually heterologous) protein. A less preferred method for the production of protein will involve transforming an unmodified host strain with a single plasmid co-expressing RNase and the recombinant protein.

RNA may be removed from a protein preparation as follows. Polyethylenimine, a polycation which binds strongly to nucleic acids to form a precipitate can be added to a clarified cell lysate containing 10mg/ml protein in 0.1M Tris-HCl, pH 8 at 4°C. The precipitate is removed by centrifugation (Brewer, S.J. and Sassenfeld, H.M. Engineering proteins for purification. In Protein purification applications: a practical approach., E. Harris and S. Angal eds., IRL Press Oxford, UK, 1990).

RNA may be removed from a cellular component such as protein using diatomaceous earth as described by Horn *et al.* (U.S. Pat. No.: 5,576,196).

Protein is purified from a host cell preparation according to a protein purification procedure specific for the protein of interest and as known in the art. Methods for protein purification are described in Protein Purification; Principles and Practice, R. Scopes, ed., C.R. Cantor, Springer-Verlag, 1985). In general, such protein purification procedures will include the steps of culturing a host cell containing the protein expression vector, inducing protein expression, and extracting and purifying the protein of interest. For example, metal chelate chromatography can be used to purify Influenza Nuclear protein that has been tagged with histidine residues (described in Example V).

Host cells other than *E. coli* can be transformed with an RNase gene and a gene of interest by the method of PEG transformation as described in PEG Transformation of Streptomyces: Wohlleben & Moth in Chapter 6 of Plasmids; A practical Approach (supra).

Host cells other than *E. coli* may also be transformed with an RNase gene and a gene of interest by the method of electroporation transformation as described in Electroporation Transformation of gram positive bacteria, Alonso & Espinosa, in Chapter 2 of Plasmids; A Practical Approach, supra).

### Regulated RNase Gene Expression

To reduce the toxic affects of RNase on the host cell, the RNase gene can be expressed under the control of an inducible promoter.

Inducible promoters that have been used to regulate the activity of RNase in *E. coli* include trp, T7, Ptac and the P<sub>R</sub> and P<sub>L</sub> promoters of phage lambda.

#### 5 Method for Inducing RNase expression in the cytoplasm

The expression of the RNase gene may be regulated by an inducible promoter that allows only minimal expression of the RNase gene under uninduced conditions so as to prevent toxic effects of RNase during the period of uninduced growth.

A variety of inducible promoter systems can be used to regulate RNase according to the method of the claimed invention. Many of the promoters (described above) can be manipulated so that expression of the RNase gene is either increased or decreased.

To inducibly express RNase in the cytoplasm RNase expression will be induced 1-3 hours prior to the harvesting of the cells producing the cellular component of interest to allow intracellular RNA degradation to occur.

15 The duration of the induction period as well as the period during which RNase is expressed will be optimized to allow maximum degradation of RNA to occur without adversely effecting product quality or cellular integrity. This will be done by incubating cells expressing RNase for 5 min to 2 hours, at 4°-37° or 42°, if induction by temperature is being employed, at a pH of 5-8 or, in the case of RNase A, a pH of 5-10.

20 For example, if the cellular component being produced is a recombinant plasmid, conditions will be modified to avoid lysing the cells as a result of over expression of intracellular RNase.

Upon completion of RNA degradation (determined as described below), cells will be lysed and the cellular component will be purified (described below).

#### 25 RNase gene under control of an inducible promoter

The toxic affects of RNase have also been reduced by expressing the RNase gene under the control of an inducible promoter. Inducible promoters that have been used to regulate the activity of the RNase gene include trp, T7, trc, lac, P<sub>tac</sub> and the P<sub>R</sub> and P<sub>L</sub> promoters of phage lambda in *E. coli* (Fujimura *et al.*, supra, Schein *et al.*, supra, Laity  
30 *et al.*, supra, Quaas *et al.*, supra, Ribo *et al.*, supra, DelCardayre *et al.*, supra, Okorokov

*et al.*, supra, McGeehan and Brenner, *et al.*, supra, Tarragona-Fiol *et al.*, supra). By using an inducible promoter to regulate the expression of the RNase gene, the level of RNase protein can be maintained at a low level under non-inducing conditions.

### 1. Trp promoter

5 A vector containing an RNase gene under the control of a *trp* promoter can be constructed according to the methods of Fujimura *et al.*, supra and Schein *et al.*, supra. Briefly, the pGH-L9 vector containing the *trp* promoter and the hGH gene, will be digested with *Cla*I and *Sal*I. The linearized vector containing the *trp* promoter will be isolated from a 1% agarose gel after electrophoresis. An RNase gene will be excised  
10 from a vector by digestion with *Cla*I and *Sal*I or amplified by PCR, using 5' and 3' RNASE specific primers that contain a *Cla*I or a *Sal*I site on the 5' end, respectively. The DNA fragment containing the RNase gene will be isolated from a 5% polyacrylamide gel after electrophoresis. The *trp* promoter-containing vector and the RNase gene will be mixed and incubated with T4 DNA ligase, and the desired plasmid  
15 will be obtained from *E. coli* HB101 transformants by standard procedures (Fujimura *et al.*, supra).

### 2. T7 promoter

The RNase gene can be cloned downstream of the T7 promoter (from  $\phi$ 10). Expression of T7 RNA polymerase will be controlled by an inducible promoter such as  
20 the lacUV5 promoter from *E. coli* BL21 (Neubauer and Hofmann, supra) or the  $\lambda$ PL promoter in conjunction with the cI857 temperature sensitive repressor (Tabor and Richardson, supra).

The lacUV5 system is based on the *lacUV5*-controlled expression of the T7-RNA polymerase. The RNase gene can be cloned behind the strong  $\phi$ 10 promoter of the T7  
25 phage, which is only transcribed by the T7-specific RNA polymerase (Neubauer and Hofmann, supra). This system can be utilized to regulate the expression of an RNase gene by the T7 promoter according to the following method of Neubauer *et al.*, Appl. Microbiol. Biotechnol., 36: 739-744, 1992.

*Escherichia coli* BL21 (F<sup>-</sup> hsdS gal omp T r<sub>B</sub><sup>-</sup> m<sub>B</sub><sup>-</sup>) carrying the lambda derivative  
30 DE3 in the chromosome DNA on which was cloned the T7 RNA polymerase gene under P<sub>lac</sub> control and the plasmid plysS. A BamHI-AluI fragment containing an

RNase gene can be cloned into the *Bam*H1-*Alu*I digested plasmid pET3, behind the  $\phi$ 10 promoter of the T7 phage.

This promoter has only a recognition site for the T7 RNA polymerase. Therefore the expression of the RNase gene should be very low under restrictive conditions.

- 5 However, after induction of the T7 RNA polymerase synthesis by IPTG (Sigma, St. Louis, Mo, USA) the RNase gene should be expressed very strongly (Neubauer *et al. supra*).

The RNase gene can be regulated by the  $\lambda$ PL promoter in conjunction with the cI857 temperature sensitive repressor according to the method described in Tabor and

- 10 Richardson, *supra*.

The expression of an RNase gene can be regulated in a system that utilizes the cloned T7 Gene 1 and a T7 RNA Polymerase Promoter. The expression system consists of two compatible plasmids, pGP1-2 and pT7-1. PGP1-2, a derivative of pACYC177, provides for expression of T7 RNA polymerase. PGP1-2 consists of gene 1 of phage T7 under  
15 the control of inducible  $\lambda P_L$  promoter, and the gene for the heat-sensitive  $\lambda$  repressor, cI857.

- 20 To express a given gene, the gene is inserted into the second plasmid, pT7-1. PT7-1 contains a T7 RNA polymerase promoter,  $\phi$ 10, isolated from a 40-bp T7 fragment. A polylinker containing eight different restriction sites lies adjacent to the promoter to facilitate the insertion of DNA fragments. Transcription from the  $\phi$ 10 promoter results in expression of the cloned gene and the  $\beta$ -lactamase gene. Exclusive expression of these genes is achieved, after heat induction of T7 RNA polymerase, by the addition of rifampicin to shut off *E. coli* RNA polymerase transcription (Tabor and Richardson, *supra*).

- 25 A vector containing an RNase gene under the control of a T7 promoter can also be constructed according to the method of Okorokov *et al.*, *supra*, Laity *et al.*, *supra*, delCardayre *et al.*, *supra* and Ribo *et al.*, *supra*.

- 30 Briefly, the expression vector pET21d(+) can be used for the insertion of an RNase A gene in conjunction with the bacteriophage DE3 lysogen strain BL21(DE3) *E. coli*, which has the T7 RNA polymerase gene under control of the *lac* UV5 promoter inserted into the chromosome. The *Bam*HI-HindIII fragment of pUCBFRA, with the RNase A

coding sequence can be recloned into the pET21d(+) vector. To keep the open reading frame, a 19-bp *NcoI-BamHI* fragment from the multiple cloning site of pTZ19RJL1 will be inserted in the corresponding unique sites. This method has been used by Okorokov *et al.* to construct the plasmid pETFRA (Okorokov *et al.*, supra). This method can also  
 5 be used to produce a vector containing other RNase genes under the control of a T7 promoter.

### 3. tac promoter

A vector containing an RNase gene under the control of a ptac promoter is constructed according to the following methods of Tarragona-Fiol *et al.* Supra.

- 10 Primers complementary to the 5' and 3' ends of the coding sequence for an RNase precursor can be synthesized and used for a PCR amplification. The 5' primer (primer 10) will contain additional sequence information which will incorporate a short ORF in front of the RNase precursor and an *EcoRI* restriction site with four additional nt to ensure correct cleavage. The 3' primer (primer 11) will contain the termination codon  
 15 for RNase, an *EcoRI* site and an additional 4 nt to ensure correct cleavage.

The two-cistron fragment resulting from PCR amplification using these primers will contain two sets of coding sequences: one for a hexapeptide and the other for an RNase precursor. Transcription of this fragment will produce bicistronic mRNA which upon translation generates a hexapeptide and the pre-RNase.

- 20 The PCR reaction will contain pQR138 as template (100 ng)/primers 10 and 11 (50 pmole/each)/all four dNTPs (0.2 mM each)/20 mM Tris-HCl pH 8.0/15 mM  $(\text{NH}_4)_2\text{SO}_4$ /2 mM  $\text{Mg}_2\text{Cl}$ /0.05%NP40/0.05% Tween 20, in a total volume of 50 or 100  $\mu\text{l}$ . The reaction components will be rapidly mixed followed by centrifugation at 10,000 x *g* (average) for 2 s at room temperature. An equal volume of paraffin oil will be  
 25 added to avoid evaporation during the reaction. Template DNA will be fully denatured by incubation for 5 min at 92°C for 1.5 min, annealing at 55°C for 1 min, and extension at 72°C for 1.5 min.

- The plasmid pQR163 contains two ribosome binding sites, one provided by the *tac* promoter of the vector and the other, for translation of the second cistron, is contained  
 30 within the coding sequence of the first cistron. The mRNA produced upon IPTG induction of *E. coli* cells harboring pQR163 is bicistronic and starts from the *tac*



promoter. The first cistron encodes a 6-amino acid peptide (Met-Phe-Leu-Glu-Asp-Asp). The stop codon of the first cistron and the start codon of the second cistron overlap such that ribosomes will continue translation of the mRNA and produce pre-RNase (Tarragona-Fiol *et al.*, supra).

- 5 According to the method of Tarragona-Fiol, the endogenous RNASE signal sequence is present in the RNase expression vector. However, other signal sequences can be cloned onto the RNASE gene, as described below.

If the endogenous signal sequence is present, the synthesized precursor form of RNase will be translocated to the periplasm. N-terminal sequencing will be used to show that  
 10 the signal sequence has been correctly cleaved. The oxidative environment of the periplasm will allow the correct folding of RNase to form the native enzyme as evidenced by the recovery of fully active enzyme (Tarragona-Fiol *et al.*, supra).

#### 4. $P_R$ and $P_L$ promoters of phage lambda

- 15 A vector containing an RNase gene under the control of the  $P_R$  promoter of phage lambda is constructed according to the following method of Okorokov *et al.* Supra.

The  $P_R$  expression cassette can be isolated from the vector pCQV2 containing the  $P_R$  promoter of the *E. coli*  $\lambda$  phage under cI repressor regulation. The cassette is surrounded by an *EcoRI* restriction site downstream of the cI coding frame and a *BamHI* site downstream of the  $P_R$  promoter.

- 20 RNase genes can be isolated from a vector by using appropriate restriction enzymes that are compatible with pCQV2. The RNase gene can be cloned under the control of the  $P_R$  promoter in the absence of a signal sequence. RNase genes can also be isolated from a vector by PCR using primers that produce an amplified fragment that contains the RNase gene and has the appropriate restriction sites on the ends.

- 25 A vector that can be used to clone the expression cassette fragment containing the RNase A gene is pUC19. The fragments *EcoRI-BamHI* from pCQV2 and *BamHI-HindIII* from pUCBFRA, containing the *phoA* signal peptide-RNase A fusion frame, can be joined with pUC19, restricted with *EcoRI/HindIII*, in a three-component ligation reaction. This method has been used by Okorokov *et al.*, supra to construct the plasmid  
 30 pEXFRA.

A vector containing an RNase gene under the control of the  $P_L$  promoter of phage lambda is constructed according to the following method of McGeehan and Breener, *supra*.

5 The gene for RNase can be cloned in vector pN1, and transferred into plasmid pAL181, together with a linker containing the methionine start codon. In this plasmid, the RNase gene will begin just after the Shine Dalgarno sequence in pAL181.

#### 5. Lac promoter

A vector containing a gene regulated by the lac operator/promoter in conjunction with the lac repressor may be constructed as described in the section entitled cloning RNase  
10 into pN1 or according to Neubauer and Hofmann, *supra*.

A vector containing an RNase gene under the control of the lac operator/promoter is constructed according to the following method of Quass *et al.* *supra*.

pIN-III-*ompA2* will be digested with *Hind*III and *Eco*RI. An RNase gene digested with the same enzymes can be inserted into pIN-III-*ompA2*. Since the entire gene is under  
15 the control of the *lpp* promoter and the *lac* promoter-operator, gene expression is regulated by the *lac* repressor, i.e. inducible by adding isopropyl- $\beta$ -D-thiogalactoside.

This method has been used by Quass *et al.* to produce the plasmid pA2T1-1 in which the RNase T<sub>1</sub> gene is fused in frame to the *ompA* signal peptide gene. However, the lac promoter can also be used to regulate the expression of an RNASE gene fused in frame  
20 to a variety of signal sequences (described below).

#### 6. Trc promoter

A vector containing an RNase gene under the control of the trc promoter is constructed according to the method of Okorokov *et al.* *supra*.

An expression vector which provides the strong inducible *trc* promoter is pKK233-2.  
25 pKK233-2 features the *lacZ* ribosome binding site and an ATG initiation codon, which is accessible by cleavage at the unique *Nco*I restriction site. The multiple cloning site is followed by the gene encoding for 5S rRNA and the strong transcription signals T1 and T2 of *rrnB E. coli* (Okorokov *et al.* *supra*). An RNase gene can be isolated from a vector by digestion with enzymes that are compatible with the multiple cloning site of  
30 pKK233-2. Alternatively, an RNase gene can be amplified by PCR using primers that

direct the synthesis of a fragment that has restriction enzyme sites that are compatible with the multiple cloning site of pKK233-2.

### Inducible promoters systems

#### 1. Temperature Inducible Promoters

- 5 The  $P_L$  promoter can be used in conjunction with the cI857I repressor to regulate the expression of the RNase gene.

A vector expressing an RNase gene under the control of the  $P_L$  promoter can be produced as described in the section entitled RNase gene under control of an inducible promoter.

#### 10 Induction

The cI857I repressor is temperature sensitive and represses gene expression at 30°C. When the temperature is increased to 42°C, the repressor is inactivated and expression occurs (Tabor and Richardson, *supra*).

- Induction can be carried out according to the following method of McGeehan *et al.*, *supra*.
- 15 LB medium (50 ml) containing ampicillin (125 µg/ml) will be inoculated with a fresh colony formed from the growth of *E. coli* transformed with a vector comprising an RNase gene under the control of the  $P_L$  promoter, and then grown to density overnight. This culture will be used to inoculate four 5L Erlenmeyer flasks each containing 1 liter of LB/amp (100 mg/ml) medium. These cultures will be grown at 30°C to an  $A_{550}$  value
- 20 of 1.2 in a shaker. The flasks will be transferred to a shaker warmed to 42°C, and LB medium (400 ml) preheated to 72°C will be added to each flask to bring the temperature of each quickly to 42°C. Shaking will be continued for 25 min, and the cells will be recovered by centrifugation (10 min at 4000 x g). The bacterial paste will be resuspended in 10 x (v/w) denaturing buffer (8 M urea, 40 mM sarcosine, 20 mM
- 25 Tris, 30 mM mercaptoethanol, 1 mM EDTA, pH 7.8). The suspension will be passed through a French press (4000 lb/inch<sup>2</sup>) and the insoluble materials removed by centrifugation for 10 min at 15000 x g. The clarified material will be dialyzed ( $M_r$  > 3500 cutoff) vs two changes of refolding buffer (100 mM NaCl, 50 mM Tris, 4 mM oxidized glutathione, 4 mM reduced glutathione, 1 mM EDTA, 0.4 mM
- 30 phenylmethylsulfonyl fluoride (PMSF), 0.02% NaN<sub>3</sub>, pH 7.6). The dilysate will be loaded onto a Sephadex G50 sizing column and eluted at a linear flow rate of 2.5-3ml

h<sup>-1</sup> cm<sup>-2</sup> using elution buffer (100 mM NaCl, 50 mM Tris, 0.4 mM PMSF, 0.02% NaN<sub>3</sub>). Fractions containing RNase will be identified by catalytic activity against a synthetic RNA (UpA). These fractions will be pooled and concentrated 4-fold using an Amicon ultrafilter device with a YM5 filter ( $M_r > 5000$  cutoff). This solution will be  
 5 dialyzed into a new buffer (100 mM NH<sub>4</sub>OAc, 0.02% NaN<sub>3</sub>, pH 6.1) and then applied to a column of affinity resin (4 ml, pUp Sepharose, Pharmacia) by gravity. The eluate under these conditions should be essentially free of RNase activity. The column will be washed with 10-20 vols of buffer, and the RNase will be eluted using a new buffer (200 mM NH<sub>4</sub>OAc, 300 mM cytidylic acid, pH 6.1). Small fractions (1.5 ml) will be  
 10 collected. Fractions containing RNase activity will be pooled and dialyzed vs two changes of buffer (100 mM NH<sub>4</sub>OAc pH 6.1) to remove the cytidylic acid. The solution will be placed in a sterile tube and stored at 4°C (McGeehan and Benner, *supra*).

A vector expressing an RNase gene under the control of the P<sub>R</sub> promoter can be  
 15 produced as described in the section entitled RNase gene under control of an inducible promoter.

### Induction

Induction can be carried out according to the following method of Okorokov *et al.*, *supra*). A 6-h culture of the *E. coli* strain carrying the expression vector comprising an  
 20 RNase gene under the control of the P<sub>R</sub> promoter, for example pEXFRA, will be grown in LB medium supplemented with ampicillin (125 µg/ml) at 30°C at 220 rpm. This culture will be used for inoculation of rich medium containing 24 g/liter yeast extract, 12g/liter tryptone, 12.54 g/liter, 12.54 g/liter K<sub>2</sub>HPO<sub>4</sub>, KH<sub>2</sub>PO<sub>4</sub>, 4ml/liter glycerol, and 125 mg/liter ampicillin and adjusted to pH 7.1.

25 After growth at 28°C and 220 rpm in rich medium to a density of OD<sub>600</sub> ~ 0.6, the culture will be induced by heat shock. The following heat shock protocols can be used. The temperature of the culture can be shifted to 42°C by adding one-half volume of prewarmed (85°C) rich medium. The culture will then be agitated at 42°C, 120 rpm, for 30 min. The temperature will then be reduced to 37°C and shaking (220 rpm) will be  
 30 continued for a further 10 to 12 h.

Alternatively, the temperature of the culture can be shifted to 37°C by inoculating directly into prewarmed (37°C) rich medium. The culture will then be grown for an additional 16-18 h at 37°C and 220 rpm.

## 2. Inducer Controlled Promoters

- 5 The lac operator/promoter in conjunction with the lac repressor can be used to regulate the expression of the RNase gene.

A vector expressing an RNase gene under the control of the lac operator/promoter can be produced as described in the section entitled RNase gene under control of an inducible promoter.

## 10 Induction

Fermentation will be carried out in a medium wherein the lac operator is repressed for example LB medium without lactose or isopropyl- $\beta$ -D-thiogalactoside (IPTG).

- Expression will be induced by the addition of IPTG to a final concentration of 0.5mM or lactose (Neubauer and Hofmann, FEMS Micro. Reviews:14, 99-102, 1994, Quass *et al.*, supra, and Donovan *et al.*, J. Industrial Microbiology, 16: 145-154, 1996).
- 15 Induction with IPTG can be carried out as described in example 3 below, and according to the method of Quass *et al.*, supra.

- To induce transcription of an RNase gene under the control of the lac promoter, IPTG will be added to a final concentration of 0.1 to 0.5 mM to a liquid culture grown to an
- 20 absorbance of 0.5 at 600nm. Cells will be allowed to grow overnight and will be harvested by centrifugation (8500 x g, 15 min, 4°C). RNase will be released from the cells by osmotic shock following the modified protocol of Koshland and Botstein. The pellet of a 1 liter culture will be resuspended in 40ml ice-cold Tris/HCl/EDTA buffer, (50 mM Tris/HCl, pH 7.5, 10 mM EDTA) containing 15% sucrose and kept on ice for
- 25 30 min. After centrifugation the cells will be resuspended in ice cold Tris/HCl/EDTA buffer, incubated for another 30 min on ice and centrifuged again. The supernatants from the two washing steps will be combined, the pH adjusted to 2.5 with 15% HCl at 4°C and the precipitated proteins separated by centrifugation (12000 x g, 25 min, 4°C). The pH of the supernatant will be adjusted to 7.5 with 10 M NaOH, diluted to 200 ml
- 30 with Tris/HCl EDTA buffer and applied to a DEAE-Sepharose CL-6B column (12 ml column volume) equilibrated with Tris/HCl/EDTA buffer. Recovery will be achieved

by gradient elution (100-700 mM NaCl in Tris/HCl/EDTA buffer). Fractions containing RNase activity will be combined and passed through a Bio-Gel P-30 column (1.5 x 50 cm) using 50 mM ammonium acetate as running buffer (Quass *et al.*, supra).

The strength of the lac promoter could be manipulated to increase expression by cloning the trp promoter upstream of the lac promoter to create the tac promoter.

The affinity of the lac repressor for its operator can be manipulated to increase or reduce binding and, subsequently, expression of the gene under the control of the lac promoter (Müller, *et al.*, J. Mol. Biol., 257: 21-29, 1996).

### 3. Starvation Inducible Promoters

To control the amount of RNase being produced in a host cell, the RNase gene can be placed under the control of the trp operator in conjunction with the trp repressor.

A vector expressing an RNase gene under the control of the trp operator in conjunction with the trp repressor can be produced as described in the section entitled RNase gene under control of an inducible promoter.

#### Induction

During growth in the presence of tryptophan, expression of the controlled gene is repressed (Schein & Noteborn, Biotechnology, 6: 291-294, 1988), and upon tryptophan starvation expression of the gene is induced. The trp promoter can also be induced by the introduction of IAA (3-indoleacrylic acid), according to the method of Fujimura *et al.*, supra.

*E. coli* harboring a plasmid comprising an RNase gene under the control of a trp promoter, for example, pTPW1, will be grown at 37°C in 4 liters of M9-casamino acid medium containing ampicillin (20 µg/ml). After incubation for 1.5h (optical density of culture at 660nm, 0.05-0.1), 3-indoleacrylic acid (IAA, 40 µg/ml) will be added to the medium to induce the expression of the fusion gene. Cells will be allowed to grow further for 8 h, harvested by centrifugation (3500 x g, 10 min, 4°C), and washed in isotonic saline (Fujimura *et al.*, supra).

The following purification steps can be used to purify an RNase protein that has been secreted to the periplasmic space. Proteins in the periplasmic space will be released by

osmotic-shock treatment by a modified version of the protocol of Cornelis *et al.* The saline washed cells will be suspended in 1 liter of 25% sucrose containing 1mM EDTA and 30 mM Tris-HCl (pH 8.0) and the suspension will be shaken at 37°C for 30 min. The cells will then be sedimented by centrifugation (14000 x g, 1 h, 20°C), resuspended in 1 liter of ice cold 20 mM Tris-HCl (pH 7.5), and shaken at 0°C for 30 min. The suspension will be centrifuged (10000 x g, 15 min, 4°C) and the resultant supernatant will be used as the periplasmic fraction. The cytoplasmic fraction will be prepared from sedimented cells by disruption with lysosome in the presence of SDS. RNase will be purified from the periplasmic fraction by two steps of column chromatography: first, anion-exchange chromatography on a Q-Sepharose column equilibrated with 0.15 M NaCl in 20 mM Tris-HCl (pH 7.5) and eluted with a linear NaCl gradient (0.15-0.45 M) in 20 mM Tris-HCl (pH 7.5); then gel chromatography on a Sephadex G-50 column from which the enzyme was eluted with 20 mM ammonium bicarbonate. The enzyme at each purification step will be analyzed by SDS-PAGE (Fujimura *et al.*, supra).

#### 4. Bacteriophage promoters

The T7 RNA polymerase/promoter system includes a promoter that is stronger than the promoter systems described above.

A vector expressing an RNase gene under the control of the T7 RNA polymerase/promoter system can be produced as described in the section entitled RNase gene under control of an inducible promoter.

An RNase gene will be cloned downstream of the T7 promoter ( $\phi 10$ ). Expression of T7 RNA polymerase will be controlled by an inducible promoter such as the lacUV5 promoter as is found in *E. coli* BL21 (Neubauer and Hofmann, supra) or  $\lambda P_L$  promoter with the cI857 temperature sensitive repressor (Tabor and Richardson, 82: 1074-1078, 1985).

Expression of the RNase gene will depend upon the expression of the bacteriophage RNA polymerase. In the event that expression of the RNase from the T7 promoter is leaky under uninduced conditions, then a transcriptional terminator for the *E. coli* RNA polymerase could be inserted into the desired promoter for the T7 RNA polymerase such that

expression of the polymerase is dependant upon transcriptional read through (Tabor and Richardson, supra).

Expression of an RNase gene under the control of the T7 promoter can be obtained by the following method of Okorokov *et al.*, supra.

- 5 A 6-h culture of the *E. coli* strain carrying an expression vector comprising an RNase gene under the control of the T7 promoter, for example pETFRA, will be grown in LB medium supplemented with ampicillin (125  $\mu$ g/ml) at 30°C at 220 rpm. This culture will be used for inoculation of rich medium containing 24 g/liter yeast extract, 12g/liter tryptone, 12.54 g/liter , 12.54 g/liter K<sub>2</sub>HPO<sub>4</sub>, KH<sub>2</sub>PO<sub>4</sub>, 4ml/liter glycerol, and 125  
10 mg/liter ampicillin and adjusted to pH 7.1. After growth at 28°C and 220 rpm in rich medium to a density of OD<sub>600</sub> ~ 0.6, expression will be induced by adding IPTG to a final concentration of 1 mM with a further incubation for 10 h (Okorokov *et al.*, supra).

#### 5. Inhibitor Induction

- The *Bacillus* RNase, BaRNase, is inactive in the presence of its specific intracellular  
15 inhibitor Barstar (Hartley, supra). The presence of the barstar gene in the host cell which expresses baRNase suppresses the lethal effect of baRNase activity when both genes are equally expressed. The presence of the barstar gene under the control of a repressible promoter could be used to express baRNase activity at the termination of fermentation.
- 20 When the gene for wild-type baRNase, which is lethal alone, is assembled on the same plasmid as the barstar gene on a pUC19-pC194 shuttle vector, baRNase is correctly processed and secreted by *Bacillus subtilis* but remains cell-bound in *E. coli*. With barstar and the *phoA* promotor-signal sequence construct, which has an *E. coli* signal  
25 peptidase site at the N terminus of mature baRNase, following induction by low phosphate, authentic baRNase is secreted by *E. coli* at a level of 20 mg/l. A variable amount of this baRNase, generally about half, will appear in the culture medium, the rest remaining in the periplasmic space. On the addition of acetic acid to 5%, however, all of the baRNase will be released to the medium, from which it may be directly absorbed onto phosphocellulose. More than 90% of the protein eluting from the  
30 phosphocellulose in high salt is baRNase. Essentially pure baRNase may then be



obtained by salt gradient chromatography on a strong anion exchanger such as SP-Trisacryl (Pharmacia) (Hartley, *supra*).

## 2. Inducible RNase expression in the periplasmic space or the cellular cytoplasm

### RNase gene secreted to periplasm or cellular supernatant

- 5 To prevent both toxic effects of overproduced RNase on the host cell, and proteolysis of the exogenous RNase by host cell proteases, the RNase gene is modified to encode a protein that is not secreted into the cytoplasm of the host cell. RNases may be produced as fusion proteins with signal peptides that direct secretion of the enzyme to either the periplasmic space or the culture supernatant. The signal peptides that have been used
- 10 successfully to direct RNase out of the cytoplasm include the endogenous RNase secretion signal (Schein *et al.*, *supra*, Tarragona-Fiol *et al.*, *supra*, Russo *et al.*, *supra*), the *E. coli* alkaline phosphatase signal peptide (phoA) (Fujimura *et al.*, *supra*, Okorokov *et al.*, *supra*), the pelB sequence (Ribo *et al.*, *supra*, DelCardayre *et al.*, *supra*), and the signal peptide of the OmpA protein of the *E. coli* outer membrane (Quaas *et al.*, *supra*).
- 15 RNase genes may also be expressed as fusion proteins. For example, an RNase A- gene 10 fusion protein is expressed as an inclusion body (Laity *et al.*, *supra*, Nambier *et al.*, *Eur. J. Biochem.* 163: 67-71, 1987).

The presence of the gene coding for the Bacteriocin Release Protein (BRP) in a host cell allows for secretion of the overexpressed RNase into the culture supernatant rather than

20 into the periplasm. When the BRP gene is moderately induced, lysis of the host cell is avoided, allowing for large-scale protein production in continuous culture (van der Wal *et al.*, *FEMS Microbiol. Rev.* 17: 381-399, 1995). This system may be used for secretion of RNase.

By directing the export of the RNase enzyme to a location outside of the host

25 cytoplasm, problems of RNase toxicity and RNase degradation by proteolysis are minimized.

Induction of RNAase expression will be regulated according to the parameters defined in the section entitled inducible RNase expression in the cytoplasm. However, RNAase will be secreted into the periplasmic space and will not contact the intracellular product

30 until cell lysis.

To target newly made RNase A out of the cytoplasm, a leader sequence can be cloned onto the RNase gene such that the protein is secreted into either the culture supernatant or the periplasmic space.

#### 1. Endogenous RNase secretion signal

- 5 The *E. coli* RNase I gene has an endogenous leader sequence which directs the gene product to the periplasm, and, when overexpressed on a high copy number plasmid does not affect cell growth (Meador and Kennell, *Gene*, 95:1-7, 1990).

The endogenous RNase leader sequence can be cloned onto an RNase gene as described in the section entitled RNase gene secreted to periplasm or cellular supernatant.

- 10 A number of signal peptides have been used successfully to direct the secretion of an RNase protein to the periplasmic space or the cellular supernatant. These include an endogenous RNase secretion signal, the *E. coli* alkaline phosphatase signal peptide (phoA) and the pelB sequence (Fujimura *et al.*, supra, Schein *et al.*, supra, DelCardayre *et al.*, supra, Okorokov *et al.*, supra, Tarragona-Fiol *et al.*, supra).

#### 15 2. Pho A

The introduction of a phoA leader sequence onto the baRNase gene which is under the control of a tac promoter has resulted in yields of up to 1g/l BaRNase (Hartley, supra). This sequence allows for efficient processing and secretion of RNase into the culture supernatant and periplasmic space of the *E. coli* host.

- 20 The Pho A sequence can be cloned onto an RNase gene according to the method of Fujimura *et al.*, supra).

- A vector containing an RNase gene under the control of a trp promoter is constructed according to the methods of Fujimura *et al.*, supra as described. The trp promoter-containing vector, the RNase gene, and the synthetic gene for the APase signal peptide  
25 will be mixed and incubated with T4 DNA ligase, and the desired plasmid, will be obtained from *E. coli* HB101 transformants by standard procedures (Fujimura *et al.*, supra). Fujimura *et al.* have used this method to construct pTPW1, a vector comprising the RNase T1 gene under the control of the trp promoter. This method can be modified such that the synthetic gene for the APase signal peptide is be cloned onto other RNase  
30 genes (including those described) under the control of either a constitutive promoter or one of the inducible promoters described above.

### 3. Pel B

The Pel B sequence has been used to direct the secretion of RNase to the periplasm in a form that is active following a solubilization step (Ribo *et al.*, supra, delCardayre *et al.*, supra).

- 5 The Pel B sequence can be cloned onto an RNase gene by the following method described in delCardayre *et al.* Plasmid pBXR (where BXR refers to bacterial expression of RNase A) will be constructed as follows. A fragment carrying the cDNA for RNase A that can be inserted between the *MscI* and *SalI* sites of pET22B(+) will be generated by PCR. The amplified fragment will be band-purified, treated with T4 DNA
- 10 polymerase to remove any overhanging bases left by *taq* polymerase, and digested with *SalI*. The resulting fragment will have the RNase A cDNA flanked on its 5' end by two CG base pairs (which form a blunt end) and on its 3' end by a *SalI* sticky-end. This fragment will then be ligated to the band-purified *MscI-SalI* fragment of the *E. coli* expression plasmid pET22B(+) by T4 DNA ligase. A *PstI* site in the Amp gene will
- 15 then be removed by site-directed mutagenesis using oligonucleotide JHH16, making the *PstI* site in the RNase A cDNA the only such site in the plasmid. The integrity of this construction will be assessed by restriction and sequence analysis.

Plasmids comprising the pel B sequence cloned onto other RNase genes can also be constructed. A fragment containing the RNase gene and the appropriate restriction

20 enzymes (e.g. sites compatible with linearized pET22B(+)) can be ligated as described.

### 4. Bacteriocin Release Protein

The presence of the gene coding for the Bacteriocin Release Protein (BRP) in the host cell allows for secretion of the over expressed RNase into the culture supernatant rather than into the periplasm. When the BRP gene is moderately induced, lysis of the host cell

25 is avoided, allowing for large-scale protein production in continuous culture (van der Wal *et al.*, FEMS Microbiol. Reviews 17,1995, 381-399).

Cells expressing both an RNase protein and BRP can be made by co-transformation of a plasmid containing the RNase gene with a compatible plasmid containing the BRP gene. Alternatively both genes can be encoded for on the same plasmid.

Expression of the BRP gene can be induced by the addition of MitomycinC (plasmid pSW1 MoBiTec GmbH) or IPTG (plasmid pJL3, MoBiTec GmbH) (See Protocols MoBiTec GmbH, Wagenstief, 5: D-37077 or Van der Wal *et al.* supra).

The secreted RNase can be purified from the supernatant as described by Tarragona-Fiol *et al.* (supra) for RNase A, or using ThioBond™ resin if the RNase is tagged as a thioredoxin fusion (Invitrogen) or using ProBond™ metal-Binding resin if the RNase is His tagged (Invitrogen).

The purified RNase can then be added to the cell lysate/supernatant (depending upon location of the desired product e.g. intracellular plasmid/protein or extracellular protein) and incubated until RNA digestion is complete. Plasmid/protein is then purified as described in the section entitled Production of Recombinant Nucleic Acid or Protein (Horn *et al.*, supra).

#### 5. Fusion proteins

The RNase gene can also be fused with a gene that allows the RNase protein to be expressed in a soluble form or as inclusion body. For example the RNase gene could be fused with the Thioredoxin gene (R&D systems, LaVallie *et al.*, BIO/Technology 11, 1993,187-193) and, following tryptophan induction, could be selectively released upon freeze/thaw of the bacterial pellet. If the Thioredoxin-RNase fusion protein is active, there is no need to release the thioredoxin by protease cleavage.

The RNase gene of choice can be cloned in frame with the thioredoxin gene in the polylinker of pTRXFUS (Invitrogen) (LaVallie *et al.*, supra). This vector is then used to transform the production host *E. coli* G1724. *E. coli* cells will be cultured to an OD550 of 0.5 and tryptophan added to a final concentration of 0.1 mg/ml, to induce fusion protein synthesis.

If the above production host is also the host for heterologous gene expression or for the production of plasmid then the cells will be incubated until such time as the RNA has been sufficiently been digested. Protein or plasmid purification can then proceed as described.

If the above production host is not the host used for heterologous gene expression or for the production of a heterologous plasmid then the RNase fusion protein can be released by cell lysis (french press) or osmotic shock as described by LaVallie *et al.* supra, or by

freeze thaw as described by Johnson & Hecht, M.H. Bio/Technology 12:1357-1359, 1994. The released protein can then be purified using a ThioBond resin as described by Invitrogen or as described by Tarragona-Fiol *et al.* Supra.

The purified RNase can then be added to the cell lysate of the host cell producing the  
5 heterologous gene or plasmid as described (see Birnboim and Doly, supra and Ausubel, Current Protocols, supra).

#### RNase gene under control of a constitutive promoter

In some methods and host cells according to the invention, it will be preferred to constitutively express the RNase gene. Thus, the RNase gene will be placed under the  
10 transcriptional control of a constitutive promoter that is operable in the selected host cell. Constitutive promoters are well-known in the art for many different types of bacteria. For example, in *E. coli*, such promoter include the glucokinase promoter of the 6-phosphogluconate dehydrogenase promoter (Neidhardt, in Salmonella typhimurium and Escherichia coli Cellular and Molecular Biology ASM, Fraenkel, D.,  
15 Chapter 12, 0.142, 1987).

#### Constitutive Periplasmic Expression of RNAase

An RNase gene can be engineered to produce protein that is constitutively expressed in the periplasm as described in the section entitled RNase gene under control of a constitutive promoter.

20 To achieve constitutive RNAase expression in the periplasm, an induction period is not required. RNase will accumulate in the periplasm as cells grow (Tarragona-Fiol *et al.*, supra).

#### RNase Digestion In Host Cells Secreting RNase Into The Culture Supernatant

If both the cellular component of interest and the RNAase is secreted into the  
25 supernatant, whether by the same cell or different cells, cell lysis should be avoided since the product of interest will be contaminated by components released from bursting/dying cells. For example, bovine RNase A is very stable and is still active in fermentation broth at pH 6.8-7.4.

The cellular components can be collected directly from the culture supernatant without  
30 releasing other cellular derived contaminants by lysing cells. Cells will be removed

from the supernatant by centrifugation for 25 min at 10,000g. The cell-free supernatant which contains both the cellular component and the RNase, will be incubated until substantially all of the RNA has been degraded.

The optimal incubation period is the time required to degrade substantially all of the RNA from a cellular component that is either a protein, nucleic acid or CHO (as defined herein). The optimal incubation period required to degrade substantially all of the RNA from a cellular component will be determined by the following steps. At appropriate time points (5 minute intervals, up to 1 hour or more if the undegraded RNA is still present, as determined by methods described below) aliquots will be removed and assayed for the presence of RNA, by the methods described below.

Following removal of RNA the cellular component can be purified as described below.

Alternatively, it is possible to directly capture the product by applying the culture medium to an expanded bed chromatography column. According to this protocol, cells and RNase will pass directly through the column while the product will bind to the chromatography medium according to the method of Hansson *et al.*, Bio/Technology, 12: 285-288, 1994 described below.

Equipment for expanded bed adsorption. A STREAMLINE™ 50 (inner diameter of 50mm) borosilicate glass column (Pharmacia Biotech, Uppsala, Sweden), equipped with a purpose-designed liquid distributor at the bottom of the column, can be used together with the STREAMLINE™ DEAE adsorbent (Pharmacia Biotech, Uppsala, Sweden), for the ion exchange adsorption in an expanded bed procedure. The adsorbent matrix consists of spherical macroporous, cross-linked agarose particles, containing crystalline quartz to increase the bead density (mean particle density = 1.2 g/ml). The ionic capacity for the diethylaminoethyl (DEAE) adsorbent is 0.13-0.21 mmol/ml. The particle size distribution ranges from approximately 100 to 300 µm with a mean particle size of 200 µm. The adsorbent is autoclavable, has an operational pH stability between 3 to 9 and resists

1M NaOH. The bed height in sedimented configuration will be 10 cm. The linear flow rate recommended to give a suitable expansion of the bed is 200-400 cm/h. The capacity of the adsorbent to bind the target protein should be investigated in packed bed column experiments prior to setting up the process.

Expanded bed adsorption. At the end of the fermentation the pH will be allowed to drop to 5.5 by removal of the base addition for pH control  $\text{NH}_3$ -regulation or NaOH-regulation. In addition, the glycerol feed will be terminated and the temperature will be raised to 60°C for 20 minutes to increase the leakage of periplasmic proteins to the culture medium. The culture, thereafter, will be cooled to room temperature. Prior to loading, the bed will be pre-expanded two to three times with loading buffer (50 mM NaCl) at a linear flow rate of 200 cm/h. The crude fermentor broth will then be applied, from bottom to top, to the expanded bed as a 1:1 mixture with loading buffer, at the same total flow rate (200 cm/h), using a two pump arrangement allowing on line mixing. When the entire fermentor content has been loaded, the expanded bed will be kept in an expanded state by increasing the loading buffer flow rate to 200 cm/h. Approximately 1/4 volume of loading buffer and thereafter 1/2 volume of washing buffer (100 mM NaCl) will be allowed to pass through the expanded bed to wash out remaining cells and loosely adsorbed proteins. The anionic adsorbent will then be sedimented by reversing the flow and lowering the adapter to the top of the packed bed, which will have been washed with washing buffer. The packed bed will then be eluted from top to bottom using 0.5 M NaCl (flow rate 100 cm/h) and 50 ml fractions will be collected.

Analysis by viable count and OD measurements of the cell removal efficiency of the expanded bed. Samples from the bed outlet will be collected during the washing procedure. For the viable count analysis the samples will be diluted to two separate dilutions (differing a 100-fold). A volume of 100  $\mu\text{l}$  of each sample will be plated on Trypiase Blood Agar Baseplates supplemented with ampicillin (0.1 g/l). The plates will be incubated at 37°C overnight and the number of colonies will be counted. For OD measurements the samples will be diluted to give A600nm values of 0.2-0.8. For lower cell contents than A600nm = 0.01, an only viable count will be performed. A cell content up to  $10^6$  cfu/ml is below the detection level of OD measurements, indicating that a visually clear culture medium could contain 100, 000 cells per ml (Hansson *et al.*, supra).

Lysis And RNase Digestion In Cells Expressing RNase In The Periplasm Or Cytoplasm

If the RNase accumulates extracellularly in the culture supernatant and the cellular component accumulates within the cell (cytoplasm or periplasm), cell lysis is required to allow the product and contaminating RNA to come into contact with extracellular RNase.

- 5 Cells expressing RNAase in the periplasm or cytoplasm can be lysed according to the methods described in Scopes, supra or LaVallie *et al.*, supra.

#### Determination Of Residual RNA

- The invention contemplates methods for producing a cellular component that is substantially RNA-free. In order to determine whether a cellular component prepared according to the inventive methods is substantially RNA-free, the following protocols may be used.

#### RNA in DNA sample

1. Agarose gel electrophoresis based methods for RNA detection

#### Samples Preparation

- 15 Each DNA sample is initially diluted to a concentration of approximately 0.2  $\mu\text{g}/\mu\text{l}$ . Samples are then mixed in a 1:1 ratio with 2x loading buffer to a final concentration of approximately 0.1  $\mu\text{g}/\mu\text{l}$ . 10X Stock Loading Buffer is 20% w/v glycerol, 1% w/v SDS, 0.25% w/v bromophenol blue in 0.1 M  $\text{Na}_2\text{EDTA}$  (pH8.0). RNA control standards are prepared by adding 1  $\mu\text{l}$  of an approximately 1 mg/ml RNA stock solution for every 50
- 20  $\mu\text{l}$  of sample to replicate samples of DNA. 10  $\mu\text{l}$  of each sample are analyzed on a 1% agarose gel along with an aliquot of a 1 kb ladder marker (Life Technologies Ltd). Agarose gels will be run in 1X TAE buffer (40mM Tris acetate, 2mM EDTA) made from a 50X Stock solution.

#### Standards for Electrophoresis

- 25 Electrophoresis standards can be prepared by the following method. Prepare a solution of RNA Type III (from Baker's Yeast (Sigma)) at a concentration of 1 mg/ml in deionized water. This solution will then be diluted 1:5 to a concentration of 0.2  $\mu\text{g}/\mu\text{l}$ , and then 1:1 with 2x loading buffer (see recipe above) to a concentration of  $\sim 0.1 \mu\text{g}/\mu\text{l}$ . Load 10  $\mu\text{l}$  of this solution onto the gel for a 1  $\mu\text{g}$  RNA standard. Additional load 10  $\mu\text{l}$
- 30 of serial dilutions of this solution prepared according to the following method.



- Add 100  $\mu$ l of the 0.1  $\mu$ g/ $\mu$ l stock solution (in loading buffer) described above to tube 1. Remove 50  $\mu$ l from tube 1 and add this sample to 50  $\mu$ l 1x loading buffer (tube 2). Mix well by pipetting. Remove 50  $\mu$ l from tube 2 and add this sample to 50  $\mu$ l of 1x loading buffer (tube 3). Continue diluting the standard solution until 12 tubes have been prepared. Load 10  $\mu$ l of each standard solution onto an agarose gel.

#### Electrophoresis conditions

Voltage : 100 V (Controlled)

Current : 400 mA (Maximum)

Power : 100 W (Maximum)

- 10 Time : 30 minutes

#### Analysis of results

The gel is stained to visualize nucleic acid and photographed on a UV transilluminator at 254 nm.

- To detect RNA, following agarose electrophoresis, the gel is stained with either ethidium bromide or sybr Green II. RNA is visualized under UV illumination. The detection limit for ethidium bromide and sybr green II staining is approximately 3.2% w/w (RNA/total nucleic acid) and 1.6% w/w (RNA/total nucleic acid) respectively.

#### Ethidium Bromide

- If the gel is to be visualized by ethidium bromide staining, add Ethidium Bromide to the agarose gel at a final concentration of 0.5  $\mu$ g/ml and to the running buffer at a concentration of 0.5  $\mu$ g/ml. Ethidium Bromide is added from a 1 mg/ml stock solution (1 tablet of ethidium bromide per 100 mls sterile water -SIGMA lot 35H0868) stored at 4°C.

- The limits of detectability can be determined by observing which standard dilution does not permit the visualization of RNA. The preceding standard load is then expressed as a percentage of the stock standard loading (1  $\mu$ g.) If no RNA is observed in the sample tracks then quote the result as "less than X %w/w," where X is the calculated sensitivity. Should a band be observed, compare its intensity with that of the standards and estimate the amount as an amount between that of the two standards between which the

sample result lies. Quote the result on a less than/more than basis. A substantially RNA-free sample, according to this method, is less than 3.2% w/v, i.e. no RNA is detected.

#### sybr green II

- 5 To prepare a SYBR Green II solution, dilute the stock solution of SYBR Green II dye (from Molecular Probes) by 1:10,000 in TBE buffer (89mM Tris base, 89mM Boric acid, 1mM EDTA, pH 8.0). This solution can be stored in a polypropylene bottle, wrapped in aluminium foil at 4°C.

- 10 Following electrophoresis, stain the gel with SYBR Green II dye, prepared as described above, for 30 minutes at 4°C in the dark.

- 15 Determine the sensitivity of the gel by observing which standard dilution equates to no observable RNA. The preceding standard load is then expressed as a percentage of the stock standard loading (1 µg.) If no RNA is observed in the sample tracks then quote the result as "less than X %w/w," where X is the calculated sensitivity. Should a band be observed compare its intensity with that of the standards and estimate between which two standards the sample result lies. Quote the result on a less than/more than basis. A substantially RNA-free sample, according to this method, is less than 3.2% w/v, and preferably less than 1.6% w/w (i.e. no RNA is detected).

Materials and Methods for staining with ethidium bromide or Sybr green:

- 20 Equipment: Pharmacia Biotech Electricity Power Supply (or similar) an Anachem Electrophoresis Units (16 sample wells/two rows minimum) and a UV Transilluminator with 254 nm illumination.

#### 2. Northern blot analysis

- 25 Northern blot analysis of an experimental sample and standards for electrophoresis (as described in the section entitled agarose gel electrophoresis based methods for RNA detection) can be performed according to the method described in Current Protocols, supra. To detect residual RNA, add Ethidium Bromide to the formaldehyde/agarose gel at a final concentration of 0.5 µg/ml and to the MOPS running buffer at a concentration of 0.5 µg/ml. Following agarose/formaldehyde electrophoresis the gel will be rinsed in distilled H<sub>2</sub>O for 30 min. RNA will then be visualized under UV illumination.
- 30

The limits of detectability can be determined by observing which standard dilution does not permit the visualization of RNA. The preceding standard load is then expressed as a percentage of the stock standard loading (1 µg.) If no RNA is observed in the sample tracks then quote the result as "less than X %w/w," where X is the calculated sensitivity.

- 5 Should a band be observed, compare its intensity with that of the standards and estimate the amount as an amount between that of the two standards between which the sample result lies. Quote the result on a less than/more than basis. A substantially RNA-free sample, according to this method, is less than 3.2% w/v, i.e. no RNA is detected.

## 10 2. RNA Detection by HPLC assay.

According to this method, residual RNA is hydrolyzed into its constituent ribonucleotides under mildly basic conditions. The ribonucleotides are treated with an alkaline phosphatase enzyme to produce ribonucleosides which may be resolved under reverse phase high performance liquid chromatography (RPHPLC). To ensure that the

15 DNA present in the sample does not interfere with the assay, a spin filtration step is done to remove the DNA from the sample prior to analysis.

The RPHPLC conditions have been optimized for sensitivity such that this method has a limit of detection in the picomolar region; or approximately 0.1 % when expressed as a percentage of the total concentration of DNA or protein .

## 20 RPHPLC conditions;

Instrument: Hewlett Packard 1100 HPLC system. Or equivalent system capable of gradient elution and detection in the UV region.

Column: Supelcosil LC-18S 25 cm \* 2.1 mm 5µm. (Or similar)

Mobile Phase: A = 10 mM NaH<sub>2</sub>PO<sub>4</sub> pH 6.0 (97.5%); Methanol (2.5%)

## 25 B = Methanol

Gradient: 0 to 20% B over 30 minutes

Run time: 45 minutes in total.

Flow rate: 0.21 ml/min

Loading: 5 µl

Detection: UV at 260 nm

#### RPHPLC Standard preparation.

To prepare standards, transfer approximately 15 mgs of each ribonucleoside to a 50 ml volumetric flask. Dissolve the ribonucleosides and bring them to volume in mobile phase A. Combine 200µl of each ribonucleoside into a 1.5ml eppendorf and perform serial two-fold dilutions, seven times. The resulting solution can be chromatographed directly using the above conditions. The Guanosine standard may require warming and sonication to aid dissolution. Calculate the standard loading for each ribonucleoside as follows (the RMM of the ribonucleoside should be used and the result should be quoted in picomoles);

$$(Wt(g) \times 0.2 \times 1000 \times 5 \times 1) / (RMM \times 0.8 \times 50 \times 1000000 \times 2^7)$$

#### RPHPLC Sample Preparation;

Prepare samples by adding 5 µl of a 3M solution of potassium hydroxide to 50µl of sample plasmid DNA. For examples, the samples can be plasmid DNA at 2 to 3 mg/ml in Water For Injection, WFI. The solution is briefly vortex mixed and then incubated at 37°C for a minimum of 16 hours. Equal volumes of 3M acetic acid and a 1M tris buffer (pH 8) are then added; followed by the addition of 10 µl of a 100 U/ml suspension of bacterial alkaline phosphatase (BAP). The resulting solution is then briefly vortex mixed and incubated at 37°C for an additional two hours. Immediately prior to analysis the samples are placed into spin filters with molecular weight cut off membranes operating at 10,000 daltons (or as appropriate) and centrifuged at 13,000 for 10 minutes. The filtrate is chromatographed without further treatment according to the conditions described above.

#### Results;

To interpret the results, integrate the peaks corresponding to cytidine, uridine, guanosine and adenosine (the peaks will elute in this order) in the sample and in the standard chromatograms. The amount of each ribonucleotide in the sample is then calculated as follows (the RMM used here is that of the monophosphate ribonucleotide, if the standard load is quoted in picomoles then the result will be given in nanograms).

The detection limit for this method is about 0.1%w/w. A substantially RNA-free sample according to this detection method is less than 1.6% and preferably less than 0.1 to 0.5%w/w.

$$(A \times C \times D \times 75)/(B \times 5)$$

5 Where;

A = Loading of standard ribonucleoside in picomoles.

B = Area under the peak due to the standard riconucleoside.

C = Area under the peak due to the sample ribonucleoside.

D = RMM of the ribonucleotide monophosphate.

10 Sum the results obtained for each ribonucleotide to give a total RNA

content. The final result is quoted as a mass and may be used to quote the result as a percentage of the DNA or protein content (%w/w), or of the solution volume (%w/v).

## 2. RNA in protein samples

15 Residual RNA can be detected in protein samples by either the agarose gel staining methods (as described) or the HPLC assay as described. The detection limit for this method is about 0.1% w/w. A substantially RNA-free sample according to htis detection method is less than 3.2% w/w, preferably les than 1.6% and preferably less than 0.1 to 0.5%w/w.

An acceptable level of RNA contamination of therapeutic plasmid DNA is <1%w/w.

20 An acceptable level of RNA contamination of therapeutic protein is <10ng/dose.

## Determination of Amount of RNase in a Cell Lysate

An advantage of the invention is that significantly less RNase protein is needed to remove substantially all of the RNA molecules in a cell lysate, and thus significantly less RNase protein is present in the lysate.

25 The amount of RNase protein present in a cell lysate may be determined by the following methods.

## RNase I assays

Three different assays for RNase I can be used to estimate enzyme in cell extracts. Assay A measures acid-soluble ultraviolet-absorbing material released from RNA. Assay B is a more sensitive and more specific modification of assay A. <sup>32</sup>P-labeled RNA is the substrate and acid-soluble radioactive material is measured. Assay C is a sensitive method for assaying RNase I in ribosome preparations. This consists of measuring the release of ultraviolet-absorbing material from ribosomes dialyzed against urea-containing buffer. All three assays reveal latent RNase I, since, in A and B, EDTA leads to activation of the enzyme, and in C, urea causes activation. Polynucleotide phosphorylase and RNase II are inhibited by EDTA and urea and are relatively inactive on RNA.

#### (i) Assay A

A 0.1 ml reaction mixture will contain, 10 μmoles potassium phosphate buffer, and 1.0 μmole EDTA, all adjusted to pH 7.0, plus the enzyme fraction. After incubation at 37°C for 40 min, 0.3 ml of cold 3% HClO<sub>4</sub> will be added and the mixture will be cooled in ice for 15 min. After centrifugation for 15 min at 2000 g in the cold, 0.1 ml of the supernatant solution will be diluted into 1.0 ml of distilled water and the OD<sub>260</sub> determined. All assays will be carried out in duplicate and values will be corrected for any acid-soluble ultraviolet-absorbing material in the enzyme fraction. Activity is expressed as μmoles of acid-soluble nucleotide/reaction mixture/hr, assuming a molar extinction coefficient of 10,000 for the nucleotides (Gesteland, J. Mol. Biol. 16: 67-84).

#### (ii) Assay B

<sup>32</sup>P-labeled RNA will be added to the reaction mixture in assay A, keeping the total substrate concentration constant. The final specific activity of the RNA will be 2000 to 5000 cts/min/0.1 mg. After incubation, precipitation and centrifugation as in A, 0.2 ml of the supernatant solution will be removed and placed on a 2-in. aluminum planchet and dried for counting. As in assay A, activity is expressed as μmoles/hr based on the specific activity of RNA in the reaction mixture (Gesteland, supra).

#### (iii) Assay C

Visking dialysis tubing (8/22) will be boiled in dilute Na<sub>3</sub>CO<sub>2</sub> for 20 min, washed thoroughly in distilled water and stored in 0.1 ml mM EDTA. Just before use, the tubing will be washed again with distilled water. All glassware will be sterilized, and

disposable plastic gloves will be used in handling the dialysis tubing in order to minimize RNase contamination. 1.0 ml (containing 30  $\mu$ moles of nucleotides) of a solution of ribosomes that had been washed 3 times in 0.01 M-magnesium acetate, 0.005 M-tris (pH 7.5) will be dialyzed against 100 ml of 4 M-urea, 0.005 M-tris (pH 7.5), 0.05 M-KCl at 4°C. At various times a sample of the dialysis buffer will be removed and the OD<sub>280</sub> measured. Results will be expressed as  $\mu$ moles of nucleotide released/min/ $\mu$ mole of nucleotides in the original ribosomes (Gesteland, supra).

RNase A assays (Suitable for all non-specific RNases)

Recombinant RNase activity on RNA agar plates.

- 10 A solution containing 2% (w/v) agar (Bacto-agar)/100 mM MES pH 6.5 0.3% (w/v) yeast RNA will be autoclaved and poured in Petri dishes. Round holes will be made in the RNA agar using a sterile cork-borer, and recombinant or commercial RNase (0.1-25  $\mu$ g) will be applied in a volume of 150  $\mu$ l. The plates will be incubated for 2-4 h at 37°C. To visualize RNase activity, 2 M HCl will be poured over the plate. A clear zone will be produced around the hole where active RNase has been introduced and RNA that had not been hydrolyzed forms a cloudy precipitate (Tarragona-Fiol *et al.*, supra).

- 20 The amount of RNase in a cell lysate can also be measured by determining the rate of hydrolysis of cytidyl-3':5'-adenosine by RNase. The hydrolysis of CpA by RNase can be carried out at room temperature in 1-ml, 1-cm path cuvetts (Hellma). The volume of the reaction will be 1 ml. Reactions containing 0.1 mM CpA in 0.1 Tris-acetate pH 6.5 will be initiated by adding a lysate containing RNase. The hydrolysis of CpA will be measured, against a blank containing 0.1 M Tris-acetate pH 6.5 and 0.1 mM CpA, by an increase in A<sub>265nm</sub> (Tarragona-Fiol *et al.*, supra).

#### EXAMPLE I

- 25 Small scale purification of a pUC18-based *E. coli* RNaseI expression plasmid from *E. coli* DH5a without RNase A treatment

- 30 The RNaseI gene was amplified from *E. coli* strain DH1 by PCR using the primers, 5'-GGTCCTGGGGTGATTATTACGGCTGTGGC-3' and 5'-GTTTAACTCACATGATGATACTGACTGTTG-3') and cloned into the TA cloning vector pCR3.1 (Invitrogen Inc.) as described.

RNase I plate assays demonstrated that the resulting vector overexpressed the RNase I gene in *E. coli* TOP10F (Figure 3) and DH5 $\alpha$  (data not shown). RNase plate assays were performed by the following method. LB agar (supplemented with kanamycin at 30 $\mu$ g/ml) was spotted with 10 $\mu$ l of overnight cultures of; 1. (Top left) TOP10F clone 1 (pCR3.1), 2. (Top right) TOP10F clone 2 (pCR3.1 RNase I), 3. (Middle left) TOP10F clone 3 (pCR3.1 RNAase I), 4. (Middle Right) TOP10F clone 4 (pCR3.1), 5. (Bottom left) TOP10F clone 5 (pCR3.1 RNAase I) and 6. (Bottom Right) TOP10F clone 6 (pCR3.1 RNAase I) and incubated at 37°C overnight. The following day the plate was overlaid with 6ml of soft agar containing baker's yeast RNA (0.3% in 11mM EDTA. After incubation at 37°C for 4 hours, the plate was developed by the addition of 10ml of 1M HCl, and photographed.

The RNaseI gene was then cloned into pUC18 downstream of the lacZ promoter such that its expression was increased upon the addition of IPTG to a final concentration of 0.5mM. Plate assays showed expression from the pUC18 construct was marginally higher than from the pCR3.1 construct in DH5 $\alpha$  following induction (Figure 4). LB agar (supplemented with ampicillin at 50 $\mu$ g/ml and IPTG 0.5mM) were spotted with 50 $\mu$ l of overnight cultures of 1. (Top) DH5 $\alpha$  (pCR3.1 RNAase I) and 2. (Bottom) DH5 $\alpha$  (pUC18 RNAase I) cultures and incubated at 37°C overnight. The following day the plate was overlaid with 6ml of soft agar containing baker's yeast RNA (0.3%) in EDTA (11mM). After incubation at 37°C for 4 hours the plate was developed by the addition of 10ml of 1M HCl, and photographed.

When plasmids were isolated by alkaline lysis from strains containing these constructs it was apparant that the overexpressed RNaseI was unable to remove RNA from these samples whereas the addition of exogenous bovine RNAaseA could remove RNA from replicate samples. Plate assays suggested that RNaseI was either not synthesised in sufficient quantities or was sensitive to the alkaline lysis procedure (Figure 5), and in particular to the denaturation step in the presence of SDS and sodium hydroxide.

LB agar (1%) plates containing baker's yeast RNA (0.3%), Tris buffer (10mM) and EDTA (11mM) were loaded with the following samples.



1. (Top Left) DH5 $\alpha$  (pUC18 RNAse I) cells resuspended in TE buffer, lysed by the addition of 0.2M NaOH and 1%SDS and neutralised by 3M potassium acetate. The supernatant from these cells was clarified by centrifugation and 60 $\mu$ l was loaded into a well in the plate,
- 5 2. (Top Right) DH5 $\alpha$  (pUC18 RNAse I) cells resuspended in TE buffer and lysed by three cycles of freezing and rapid thawing. The supernatant from these cells was clarified by centrifugation and 20 $\mu$ l was loaded into a well in the plate,
3. (Bottom Left) DH5 $\alpha$  (pUC18 RNAse I) cells resuspended in TE buffer and lysed
- 10 sonication. The supernatant from these cells was clarified by centrifugation and 20 $\mu$ l was loaded into a well in the plate and
4. (Bottom Right) 2 $\mu$ g of bovine RNAse A in 20 $\mu$ l of TE buffer loaded into a well in the plate.
- 15 The plate was incubated at 37°C for 4 hours and then developed by the addition of 10ml of 1M HCl before being photographed.

When the protocol for small scale purification of recombinant plasmid DNA from RNaseI expressing strains was modified to include an extended incubation period (Birnboim and Doly, 1979) overexpressed RNaseI removed a portion of the

20 contaminating RNA from the preparation of plasmid DNA (Figure 6).

Specifically, the alkaline lysis procedure was modified to include a period of incubation following the neutralization step or following release of RNaseI by sonication/freeze thaw and prior to denaturation with sodium hydroxide and SDS. During this incubation period RNA was digested by the expressed RNase.

- 25 Figure 6 is an Ethidium bromide stained 0.8% agarose gel loaded with 5 $\mu$ l of the following samples prepared as described:

Lane 1. DNA fragment size marker 1 BstEII

Lane 2. DH5 $\alpha$  (pUC18) cells resuspended in TE buffer, lysed by the addition of 0.2M NaOH and 1%SDS, neutralised with 3M potassium acetate and incubated on ice for 2

hours. The supernatant was clarified by centrifugation, ethanol precipitated and resuspended in 20µl of TE buffer.

Lane 3. DH5α (pUC18) cells resuspended in TE buffer and lysed by 3 cycles of freezing and rapid thawing and incubated on ice for 2 hours. The sample was then  
5 treated with 0.2M NaOH and 1%SDS, neutralised by 3M potassium acetate and clarified by centrifugation. DNA was ethanol precipitated and resuspended in 20µl of TE buffer.

Lane 4. DH5α (pUC18) cells resuspended in TE buffer and lysed by sonication. After incubation on ice for 2 hours, the sample was treated with 0.2M NaOH and 1%SDS,  
10 neutralised by 3M potassium acetate and clarified by centrifugation. DNA was ethanol precipitated and resuspended in 20µl of TE buffer.

Lane 5. empty

Lane 6. DH5α (pUC18 RNAase I) cells resuspended in TE buffer, lysed by the addition of 0.2M NaOH and 1%SDS and neutralised by 3M potassium acetate and incubated on  
15 ice for 2 hours. The supernatant was clarified by centrifugation, ethanol precipitated and resuspended in 20µl of TE buffer.

Lane 7. DH5α (pUC18 RNAase I) cells resuspended in TE buffer, lysed by 3 cycles of freezing and rapid thawing and incubated on ice for 2 hours. The sample was then treated with 0.2M NaOH and 1%SDS, neutralised by 3M potassium acetate and  
20 clarified by centrifugation. DNA was ethanol precipitated and resuspended in 20µl of TE buffer.

Lane 8. DH5α (pUC18 RNAase I) cells resuspended in TE buffer and lysed by sonication. After incubation on ice for 2 hours, the sample was treated with 0.2M NaOH and 1%SDS, neutralised by 3M potassium acetate and clarified by centrifugation. DNA  
25 was ethanol precipitated and resuspended in 20µl of TE buffer.

## EXAMPLE II

### Use of *E. coli* DH1Rnase A (constitutively expressing RNase A) for production of plasmid pUC19tet and protein

An RNase A gene can be cloned under the control of the constitutively active  
30 glucokinase promoter and inserted into the chromosome of the *E. coli* host strain DH1

using pN1 as an intermediate plasmid (as described in the section entitled Integration of an RNase gene into the chromosome). RNase A protein will be directed to the periplasm by its endogenous leader sequence.

Plasmid production in DH1RNase A *E. coli* cells transformed with the plasmid of interest (for example, pUC19tet $\Delta$ Amp or pTX0161 described in WO 97/29190), which also expresses a chromosomal RNase A gene under the control of the constitutively expressed glucokinase promoter, grown and lysed by alkaline lysis, as described above in the section entitled Purification of Recombinant Nucleic Acid or Protein, will be performed according to the method described in WO 97/29190. The lysis protocol will be modified so that it does not include a step in which the sample is incubated with exogenously added RNase A but instead is incubated with its autologously produced recombinant RNase A.

DH1RNase A *E. coli* cells will be transformed with a plasmid containing the gene for Influenza Nuclear Protein (NP) cloned in frame into the expression vector pTRCHis (Invitrogen). The cells can be cultured as described above in the section entitled Purification of Recombinant Nucleic Acid or Protein according to the method of Horn *et al.*, supra. The cells will constitutively synthesize influenza virus nuclear protein (NP). The cells will lysed using a cell homogenizer and incubated for a length of time sufficient to produce a substantially RNA-free recombinant protein, according to the invention. The presence of residual RNA will be detected according to the methods described in the section entitled detection of residual RNA. The NP protein will be purified according to the following method.

#### Extraction and Purification

Cells will be harvested by centrifugation, resuspended in 1% lysis buffer (50mM NaPO<sub>4</sub>, 1M NaCl, 0.5% sarkosyl) and frozen at -80°C. Cells will be thawed on ice in the presence of protease inhibitors (Boehringer Mannheim complete EDTA free protease inhibitor cocktail tablets, 1 tablet/50ml of cell resuspension). DNase I will be added to a final concentration of 5mg/ml. The cell pellet will be lysed by homogenization at 30Kpsi using a single pass in a Constant Systems Ltd homogenizer. Alternative homogenization methods can be found in Scopes, supra. The lysate will be subjected to centrifugation at 10,000g to remove cell debris. Sarkosyl will be added to a final concentration of 0.5% to aid protein solubility and endotoxin removal. Following

the addition of 60 ml of 50% Ni-NTA agarose (Quiagen Inc.), the lysate will be incubated for 1 hour at 4°C to allow batch binding of the protein to occur.

The bulk of the lysate will be removed from the resin by centrifugation at 200g. The resin will be resuspended in the remaining lysate and used to pour a Pharmacia XK 26 column. The column will be washed with 400 ml Wash Buffer (50mM NaPO<sub>4</sub>, 1M NaCl, 0.5% sarkosyl, 20mM imidazole, 10% glycerol). Contaminants will be eluted with a gradient of 0% to 22% 0.5M imidazole in wash buffer. The recombinant NP (rNP) will be eluted with a step gradient of 100% of 0.5% imidazole in wash buffer using a fraction collector.

- 10 The appropriate fraction will be analyzed by SDS-PAGE to verify the presence of rNP. The fractions containing the highest concentration of rNP will be pooled (pool 1). Additionally, the fractions containing the lowest concentration of rNP will be pooled (pool 2).

- 15 Pooled rNP will be dialyzed into 25mM Hepes, 0.5% sarkosyl. The protein concentration will be determined by using a BIORAD protein assay. The presence of residual endotoxin will be determined by KQCL assay (Biowhittaker). The presence of residual RNA will be determined according to the methods described in the section entitled Determination of Residual RNA.

### EXAMPLE III

- 20 Use of *E. coli* DH1RNase A (inducible and periplasmic) for the production of plasmid pUC18 and protein

- An RNase A gene can be cloned under the control of the inducible lac promoter and stably integrated into the chromosome of DH1, using pN1 as an intermediate plasmid, according to the methods described above. RNase A protein will be directed to the periplasm by its endogenous leader sequence.

- 25 The *E. coli* cells are then transformed with the plasmid vector to be purified (for example, pUC19tetΔAmp or pTX0161) grown according to the methods described above in the section entitled Purification of Recombinant Nucleic Acid or Protein, according to the method described in WO 97/29190. RNase A production will be induced by the addition of 0.5mM IPTG prior to the termination of fermentation. Cells

will be lysed by the modified alkaline lysis procedure described in example II and the plasmid purified by the procedure described in WO 97/29190..

*E. coli* cells transformed with a vector comprising the NP gene cloned in frame into the expression vector pTRCHis (Invitrogen) and can be cultured as described above in the section entitled Purification of Recombinant Nucleic Acid or Protein according to the method described in WO 97/29190. The cells can be induced to synthesize RNase and NP protein simultaneously by the addition of 0.5mM IPTG as described above, lysed using a cell homogenizer and incubated for 10 to 60 minutes to remove RNA. The NP protein will be purified according to the methods described above (Example II and Scopes, supra).

#### EXAMPLE IV

##### Use of *E. coli* BL21(DE3)RNase A (inducible and periplasmic) for the production of plasmid and protein

An RNase A gene can be cloned under the control of the inducible T7 promoter and stably integrated into the chromosome of BL21 (DE3), using pN1 as an intermediate plasmid, according to the methods described above. RNase A protein will be directed to the periplasm by its endogenous leader sequence.

*E. coli* cells transformed with the plasmid of interest (for example, pUC19tet $\Delta$ Amp or pTX0161) and carrying the chromosomal expression cassette BL21 (DE3) RNase A comprising an RNase A gene under the control of the T7 promoter can be grown in shake flasks in LB medium. RNase A production will be induced as described in Example V below or according to the methods described in the section entitled Inducible Promoter Systems. Cells will be lysed by the modified alkaline lysis procedure described in example II and the plasmid purified as described in WO 97/29190.

*E. coli* cells BL21 (DE3) RNase A can be transformed with a vector comprising the NP gene cloned in frame into the expression vector pET (under the control of the T7 promoter). This vector is called pTX030 and is described in Example V below. The vector is cultured as described below. The cells can be induced to synthesize NP as described below, lysed using a cell homogenizer and incubated for 10 to 60 minutes to

remove RNA. The NP protein will be purified according to the methods described above (Example II and Scopes, supra).

#### EXAMPLE V

##### RNA free Influenza Virus Nuclear Protein Production in *E. coli* BL21(DE3) RNase A

- 5 *E. coli* BL21(DE3) RNase A can be produced according to the methods described for the production of DH1RNase A, using the target strain BL21(DE3) to generate BL21(DE3)RNase A in the P1 transduction step.

*E. coli* cells containing the vector BL21(DE3)RNase A will be transformed with pTX0330 and plated out on LB agar supplemented with kanamycin (LBkan) (30mg/L).

- 10 pTX0330 is a vector for T7 polymerase promoter driven expression of 'flu' NP protein tagged with six N-terminal histidine residues, generated by cloning NP coding sequence into pET-28a(+)

(Novgen Inc.). The N-terminal tag allows for purification of NP by metal chelate chromatography. PTX0330 carries the kanamycin resistance gene *kan*.

- 15 Single colonies will be used to inoculate 25ml LB supplemented with 30 µg/ml kanamycin (LB *kan*), and the resulting culture will be grown overnight at 37°C at 200 rpm. An aliquot of this culture will be used to inoculate 200 ml LB*Kan*. This subculture will be grown to an OD<sub>600nm</sub> of 1 unit. Six 1.5L aliquots of LB*Kan* will be inoculated with 30ml of this 200 ml subculture. Protein expression will be induced by
- 20 the addition of 0.5mM IPTG (final concentration) for 5 hours. Cells will then be harvested by centrifugation, resuspended in 1% lysis buffer (50mM NaPO<sub>4</sub>, 1M NaCl, 0.5% sarkosyl) and frozen at -80°C.

##### Extraction and Purification

Cells will be thawed on ice in the presence of protease inhibitors (Boehringer Mannheim complete EDTA free protease inhibitor cocktail tablets, 1 tablet/50ml of cell resuspension). DNase I will be added to a final concentration of 5mg/ml. The cell pellet will be lysed by homogenization at 30Kpsi using a single pass in a Constant Systems Ltd homogenizer. Alternative homogenization methods can be found in Scopes, supra. The lysate will be subjected to centrifugation at 10,000g to remove cell

- 30 debris. Sarkosyl will be added to a final concentration of 0.5% to aid protein solubility

and endotoxin removal. Following the addition of 60 ml of 50% Ni-NTA agarose (Qiagen Inc.), the lysate will be incubated for 1 hour at 4°C to allow batch binding of the protein to occur.

5 The bulk of the lysate will be removed from the resin by centrifugation at 200g. The resin will be resuspended in the remaining lysate and used to pour a Pharmacia XK 26 column. The column will be washed with 400 ml Wash Buffer (50mM NaPO<sub>4</sub>, 1M NaCl, 0.5% sarkosyl, 20mM imidazole, 10% glycerol). Contaminants will be eluted with a gradient of 0% to 22% 0.5M imidazole in wash buffer. The recombinant NP (rNP) will be eluted with a step gradient of 100% of 0.5% imidazole in wash buffer  
10 using a fraction collector.

The appropriate fraction will be analyzed by SDS-PAGE to verify the presence of rNP. The fractions containing the highest concentration of rNP will be pooled (pool 1). Additionally, the fractions containing the lowest concentration of rNP will be pooled (pool 2).

15 Pooled rNP will be dialyzed into 25mM Hepes, 0.5% sarkosyl. The protein concentration will be determined by using a BIORAD protein assay. The presence of residual endotoxin will be determined by KQCL assay (Biowhittaker). The presence of residual RNA will be determined according to the methods described in the section entitled Determination of Residual RNA.

## 20 Cellular Components Useful According to the Invention

### Proteins Useful According to the Invention

Proteins useful according to the methods of the invention include but are not limited to proteins that are useful according to the invention, such as receptors, enzymes, ligands, regulatory factors, and structural proteins. Therapeutic proteins also include nuclear  
25 proteins, cytoplasmic proteins, mitochondrial proteins, secreted proteins, plasmalemma-associated proteins, serum proteins, viral antigens, bacterial antigens, protozoal antigens and parasitic antigens.

Therapeutic proteins useful according to the invention also include lipoproteins, glycoproteins and phosphoproteins. Proteins or polypeptides which can be expressed  
30 using the methods of the present invention include hormones, growth factors, neurotransmitters, enzymes, clotting factors, apolipoproteins, receptors, drugs,

oncogenes, tumor antigens, tumor suppressors, structural proteins, viral antigens, parasitic antigens and bacterial antigens. Specific examples of these compounds include proinsulin, growth hormone, dystrophin, androgen receptors, insulin-like growth factor I, insulin-like growth factor II, insulin-like growth factor binding proteins, 5 epidermal growth factor TGF- $\alpha$ , TGF- $\beta$ , PDGF, angiogenesis factors (acidic fibroblast growth factor, basic fibroblast growth factor and angiogenin), matrix proteins (Type IV collagen, Type VII collagen, laminin), phenylalanine hydroxylase, tyrosine hydroxylase, oncogenes (ras, fos, myc, erb, src, sis, jun), E6 or E7 transforming sequence, p53 protein, Rb gene product, cytokine receptor, IL-1, IL-6, IL-8, viral capsid 10 protein, and proteins from viral, bacterial and parasitic organisms which can be used to induce an immunologic response, and other proteins of useful significance in the body.

The compounds which can be incorporated are only limited by the availability of the nucleic acid sequence for the protein or polypeptide to be incorporated. One skilled in the art will readily recognize that as more proteins and polypeptides become identified 15 they can be integrated into the non-viral vector of choice, transfected according to the method of the present invention and expressed in a mammalian tissue, including human tissue.

#### DNA Sequences Useful According to the Invention

##### 1. Genes Encoding Toxins

20 Examples of genes useful in the invention include those encoding such agents including but not limited to genes encoding diphtheria toxin, *Pseudomonas* exotoxin, cholera toxin, pertussis toxin, etc., as follows. Diphtheria toxin-IL2 fusions for inhibition of HIV-1 infection (Zhang *et al.*, 1992, Jour. Acquired Immune Deficiency Syndrome 5:1181); Diphtheria toxin A chain for inhibition of HIV viral production (Harrison *et al.*, 1992, AIDS Res. Hum. Retro. 8:39 and Curel *et al.*, 1993, Hum. Gene Ther. 4:71); 25 Diphtheria toxin A chain-liposome complexes for suppression of bovine leukemia virus infection (Kakidani *et al.*, 1993, Microbiol. Immunol. 37:713); Diphtheria Toxin A chain gene coupled with immunoglobulin enhancers and promoters for B-cell toxicity (Maxwell *et al.*, Cancer Res., 1991, 51:4299); Tat- and Rev- activated expression of a 30 diphtheria toxin A gene (Harrison, 1991, Hum. Gene Ther. 2:53); Diphtheria toxin-CD4 fusion for killing of HIV-infected cells (Auilo *et al.*, 1992, Eur. Mol. Biol. Org. Jour. 11:575).



Other toxins which are useful according to the invention include but are not limited to the following. Conditionally toxic retroviruses are disclosed in Brady *et al.*, 1994, Proc. Nat. Aca. Sci. 91:365 and in Caruso *et al.*, 1992, Bone Marrow Transplant, 9:187. Toxins against EBV infection are disclosed in Harris *et al.*, 1991, Cell. Immunol. 134:85, and against poliovirus in Rodriguez *et al.*, 1992, Jour. Virol. 66:1971. Toxins against influenza virus are disclosed in Bron *et al.*, 1994, Biochemistry 33:9110.

## 2. Genes Encoding Immunoactive Agents

Another agent useful according to the invention includes immunoactive agents, i.e., agents which combat viral infections or production by activating an immune response to the virus. Such agents include but are not limited to cytokines against viruses in general (Biron, 1994, Curr. Opin. Immunol. 6:530); soluble CD4 against SIV (Watanabe *et al.*, 1991, Proc. Nat. Aca. Sci. 88:126); CD4-immunoglobulin fusions against HIV-1 and SIV (Langner *et al.*, 1993, Arch. Virol. 130:157); CD4(81-92)-based peptide derivatives against HIV infection (Rausch *et al.*, 1992, Biochem. Pharmacol. 43:1785); lympho- cytotoxic antibodies against HIV infection (Szabo *et al.*, 1992, Acta. Virol. 38:392); IL- 2 against HIV infection (Bell *et al.*, 1992, Clin Exp. Immunol. 90:6; and anti-T cell receptor antibodies against viruses in general (Newell *et al.*, 1991, Ann. N.Y. Aca. Sci. 636:279).

## 3. Genes Encoding Anti-Viral Drugs

Another anti-viral agent useful according to the invention includes drugs having anti-viral activity and which are the direct product of a gene or are a product of a gene encoding a precursor of the drug, the drug then being synthesized by a biosynthetic pathway in the cell. Targets of drug intervention in the replicative cycle of, for example, a retrovirus, include (i) binding and entry, (2) reverse transcriptase, (3) transcription and translation, and (4) viral maturation and budding. Representative inhibitors of viral binding and entry for HIV include recombinant soluble CD4, immunoadhesions, peptide T, and hypericin. Nucleoside reverse transcriptase inhibitors include zidovudine, didanosine, zalcitabine, and starudine. Foscarnet, tetrahydroimidazobenzodiazepinethione compounds, and nevirapine are some non- nucleoside reverse transcriptase inhibitors. Inhibitors of transcription and translation include antagonists of the TAT gene and GLQ223. Castanospermine and protease inhibitors interfere with viral budding and maturation. Such drugs include but are not

limited to nucleoside or nucleotide analogs and products of a cellular biosynthetic pathway such as described in Harrell *et al.*, 1994, Drug Metab. Dispos. 22:124 (deoxy-guanine); Fillon *et al.*, 1993, Clin. Invest. Med. 16:339 (dauno-rubicin); Ohrvi *et al.*, 1990, Nucleic Acids Symp. 26:93 (anti-viral nucleosides); Hudson *et al.*, 1993, Photochem. Photobiol. 57:675 (thiarubines); Salhany *et al.*, 1993, Jour. Biol. Chem. 268:7643 (pyridoxal 5'-phosphate); Damaso *et al.*, 1994, Arch. Viral. 134:303 (cyclosporin A); Gallicchio *et al.*, 1993, Int. Jour. Immunol. 15:263 (dideoxynucleoside drugs); and Fiore *et al.*, 1990, Biol. Soc. Ital. Biol. Sper. 66:601 (AZT).

#### 4. Genes Encoding Proteins or Enzymes Useful for the Treatment of Patients by Gene Therapy

Genes encoding proteins or enzymes useful for the treatment of patients by gene therapy include genes encoding  $\beta$ -glucocerebrosidase for Gaucher's disease, Factor XIII and IX for hemophilia and prodrug activating enzymes for use in prodrug treatments (e.g. bacterial nitroreductases for use in gene directed enzyme prodrug therapy (GDEPT)).

#### 15 Vectors Useful According to the Invention

Vectors useful according to the invention include a vector that possesses the following characteristics:

##### i) High copy number bacterial origin of replication.

Vectors having relatively high copy number, i.e., in the range of 20-40 copies/cell up to 1000-2000 copies/cell, are especially useful according to the invention. For example, a vector that includes the pUC origin of replication is preferred according to the method of the invention. The pUC origin of replication permits more efficient replication of plasmid DNA and results in a tenfold increase in plasmid copy number/cell over, e.g., a pBR322 origin. The resulting high copy number greatly increases the ratio of plasmid DNA to chromosomal DNA, RNA, cellular proteins and co-factors, improves plasmid yield, and facilitates easier downstream purification.

##### ii) Small and stable vector backbone.

It is preferred according to the invention that the backbone of a vector used according to the methods described herein be small, i.e., less than 5 kb, and preferably 1-3 kb. The term "vector backbone" refers to the bacterial DNA necessary to maintain and propagate the vector in a bacterial host. Vectors of the invention which include both backbone and

insert will be on the order of 15-50 kb in size, or even larger. Thus, a vector backbone useful in the invention will be capable of carrying inserts of approximately 10-50 kb or larger. The insert may include DNA from any organism, but will preferably be of mammalian origin, and may include, in addition to a gene encoding a therapeutic protein, regulatory sequences such as promoters, poly adenylation sequences, enhancers, locus control regions, etc. The gene encoding a therapeutic protein may be of genomic origin, and therefore contain exons and introns as reflected in its genomic organization, or it may be derived from complementary DNA.

The vector should also be stably inherited; that is, the vector backbone preferably contains no intrinsically unstable elements prone to rearrangement, deletion, etc, such as transposons, and is stably inherited in the presence of the selective agent.

Useful vectors according to the invention include pEA $\beta$ Glu, pUC18/19tet $\Delta$ Amp, pTX0161, pUC19tet, pGL2RSV, pGL2RSVluc, pAI6tet and pCD2tatRZfull.

iii) Polylinker suitable for the insertion of therapeutic genes and regulatory sequences.

Vectors useful according to the invention include a polylinker comprising a variety of restriction sites that are useful in cleaving the vector and incorporating therapeutic genes.

iv) Absence of other bacterial protein genes.

It is preferred according to the invention that no other bacterial genes are carried on the vector backbone. Absence of other bacterial genes minimizes the possibility of a patient developing an immune response to a foreign gene or its encoded product, where the gene is present and/or expressed in the patient's cells which have been targeted with the therapeutic vector. Other bacterial genes expressed by host strain during fermentation results in a metabolic burden in the host and reduces the biomass and plasmid yields.

v) Selective marker gene.

Vectors useful according to the invention may include a gene encoding a selectable marker, e.g., an antibiotic resistance gene such as the bacterial tetracycline resistance gene. Incorporation of the tetracycline resistance gene permits the use of tetracycline as a selective agent in the plasmid preparation procedure according to the invention. One advantage to the use of a tetracycline resistance gene is that tetracycline is not degraded in *E. coli*, and therefore more tetracycline does not have to be added during

fermentation. In addition, the tetracycline resistance gene is preferred over a gene encoding ampicillin resistance because tetracycline is prescribed less often as an antibiotic in a clinical setting, and adverse responses to tet are less frequent than for amp and other  $\beta$ -lactam antibiotics.

## 5 Dosage, Mode of Administration and Pharmaceutical Formulations

The invention encompasses methods of removing RNA from preparations of cellular components.

A pharmaceutical composition comprised of a cellular component produced from a host cell that also produces an RNase may be prepared from preferably  $10^5$ - $10^8$  host cells and more preferably from  $10^6$ - $10^7$  host cells. A pharmaceutical composition comprised of a cellular component produced from a host cell that has been co-cultured with a different host cell that produces an RNase may be prepared from preferably  $10^5$ - $10^8$  of each respective host cell and more preferably from  $10^6$ - $10^7$  of each host cell. In the latter embodiment the two host cells are preferably present in the culture in a 1:1 ratio or are mixed at the point of lysis in a 1:1 ratio.

Cellular components described herein may be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in or suspension in, liquid prior to infection can also be prepared. The preparation can also be emulsified, or the cellular component encapsulated in liposomes. The term "pharmaceutically acceptable carrier" refers to a carrier that does not cause an allergic reaction or other untoward effect in subjects to whom it is administered. Suitable pharmaceutically acceptable carriers include, for example, one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the effectiveness of the vaccine. Examples of adjuvants which may be effective include but are not limited to: aluminum hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine (CGP 11637, referred to as nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-s-n-glycero-3-hydroxyphosphoryloxy)-ethylamine (COP) 19835A, referred to as MTP-PE), and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton

(MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion. Other examples of adjuvants include DDA (dimethyldioctadecylammonium bromide), Freund's complete and incomplete adjuvants and QuilA.

Cellular components of the invention can be administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories, and in some cases, oral formulations or formulations suitable for distribution as aerosols. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1%-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10%-95% of active ingredient, preferably 25-70%.

#### Other Embodiments

Other embodiments are within the following claims.

Claims

1. A method of preparing a substantially RNA-free cellular component, comprising culturing cells producing the cellular component in a medium and lysing said cells to produce a cell lysate, wherein said cell lysate contains said cellular component and  
5 sufficient RNase activity to degrade substantially all of the RNA molecules present in said cell lysate.
2. The method of claim 1, wherein the RNase is produced by said cells producing the cellular component.
3. The method of claim 1, wherein the RNase is produced by said cells in the  
10 medium other than said cells producing the cellular component.
4. A method of preparing a substantially RNA-free cellular component, comprising culturing and lysing cells producing the cellular component and cells producing an RNase in an amount sufficient to degrade substantially all of the RNA present in the preparation.
- 15 5. The method of claim 4, wherein the cells producing the cellular component also produce the RNase.
6. The method of claim 4, wherein the cellular component and the RNase are not produced by the same cells.
7. The method of claim 1 or 4, wherein said cellular component is one of a  
20 recombinant DNA, a recombinant protein, and a recombinant carbohydrate.
8. The method of claim 1 or 4, wherein the gene encoding said RNase is integrated into the genome of the cell producing the RNase.
9. The method of claim 1 or 4, wherein said RNase is non-specific.
10. The method of claim 9, wherein said non-specific RNase is RNase A, RNase M  
25 or RNase I.
11. The method of claim 1 or 4, wherein said cell producing said RNase produces said RNase in a regulated manner.
12. The method of claim 11, wherein said RNase produced in said regulated manner is overproduced by said cell producing said RNase.

13. The method of claim 11, wherein said RNase produced in said regulated manner is inducibly produced by said cell producing said RNase.
14. The method of claim 11, wherein said RNase produced in said regulated manner is constitutively produced by said cell producing said RNase.
- 5 15. The method of claim 11, wherein said RNase produced in said regulated manner is secreted out of the cytoplasm of the cell producing said RNase.
16. The method of claim 15, wherein said RNase is secreted into the periplasm of the cell producing the RNase.
17. The method of claim 16, wherein the cell is contained in a medium and said  
10 RNase is secreted out of the cell into said medium.
18. The method of claim 11, said RNase being a non-specific RNase.
19. A host cell that produces a recombinant DNA, a recombinant protein, or a recombinant carbohydrate and also produces an RNase in a regulated manner.
20. The host cell of claim 19, wherein said RNase produced in said regulated  
15 manner is overproduced.
21. The host cell of claim 19, wherein said RNase produced in said regulated manner is inducibly produced.
22. The host cell of claim 19, wherein said RNase produced in said regulated manner is constitutively produced.
- 20 23. The host cell of claim 19, wherein said RNase produced in said regulated manner is secreted out of the host cell cytoplasm.
24. The host cell of claim 23, wherein said RNase is secreted into the host cell periplasm.
25. The host cell of claim 23, wherein when the host cell is contained in a medium,  
25 said RNase is secreted out of the host cell into said medium.
26. The host cell of claim 19, said RNase being a non-specific RNase.
27. A composition comprising a host cell that produces a recombinant DNA, a recombinant protein, or a recombinant carbohydrate and a host cell that produces an RNase in a regulated manner.

28. The composition of claim 27, wherein said RNase produced in said regulated manner is overproduced.

29. The composition of claim 27, wherein said RNase produced in said regulated manner is inducibly produced.

5 30. The composition of claim 27, wherein said RNase produced in said regulated manner is constitutively produced.

31. The composition of claim 27, wherein said RNase produced in said regulated manner is secreted out of the host cell cytoplasm.

10 32. The composition of claim 31, wherein said RNase is secreted into the host cell periplasm.

33. The composition of claim 31, wherein when the host cell is contained in a medium, said RNase is secreted out of the host cell into said medium.

34. A pharmaceutical composition comprising a cellular component that is substantially RNA-free, in a pharmaceutically acceptable carrier.


15 35. A pharmaceutical composition comprising a cellular component that is substantially RNA-free obtainable by the method of claim 1 or claim 4, in a pharmaceutically acceptable carrier.



Abstract of the Disclosure


The invention relates to host cells and methods of preparing a substantially RNA-free cellular component, comprising culturing cells producing the cellular component in a medium and lysing the cells to produce a cell lysate, wherein the cell lysate contains the  
5 cellular component and sufficient RNase activity to degrade substantially all of the RNA molecules present in the cell lysate. The invention also relates to substantially RNA-free cellular components.








**FIG.2**  
INSERTIONAL MUTAGENESIS IN *E. coli*






3/4

**FIG.3**



**FIG.4**






6

1

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220



**FIG.5**



**FIG.6**



